

Chapter 1 Measurement, page 2

1.1 Imperial Measures of Length, page 11

3. Answers may vary. For example:
- a) Foot b) Inch
 - c) Foot d) Inch
 - e) Mile
4. a) Inch
5. Answers may vary. For example:
- a) Foot
7. a) 36 in. b) 189 ft.
c) 4 ft.
8. a) 10 560 ft. b) 15 yd. 2 ft. 10 in.
c) 1 mi. 703 yd. 1 ft.
9. 165 in. = 4 yd. 1 ft. 9 in.
10. a) 52 ft. = 17 yd. 1 ft. b) \$197.82
11. a) 24 mats
12. No; 21 ft. 9 in. = 7 yd. 9 in.
13. 10 in.
14. a) 39 ft. 2 in. b) 4 rolls
c) \$49.96
15. a) \$119.99 b) \$18.59
16. 1062 ft.
17. 62 mi.
18. 28 tulip bulbs
19. 2 mi. 80 yd.
20. 1:2 349 000
21. a) \$351 000
22. \$158 400 000

1.2 Math Lab: Measuring Length and Distance, page 15

3. Calipers require a steady hand to ensure an accurate reading. Calipers cannot be used for large measures.

1.3 Relating SI and Imperial Units, page 22

Answers will vary depending on the conversion ratios used.

- 4. a) 40.6 cm b) 1.2 m
c) 4.6 m d) 1.5 km
e) 9.7 km f) 50.8 mm
- 5. a) 1 in. b) 8 ft.
c) 11 yd. d) 93 mi.
- 6. a) 55.9 cm b) 256.5 cm
c) 9.6 m


- 7. a) i) 2 ft. 6 in. ii) 3 yd.
 iii) 6 mi.
- 8. 100.6 m by 54.9 m
- 9. Tennessee River
- 10. The odometer is accurate; 142 km is close to 87 mi.
- 11. a) The warehouse
- 12. a) Michael
- 13. a) CN Tower: approximately 1815 ft.;
 Willis Tower: approximately 442.3 m
b) CN Tower c) 111 m; 364 ft.
- 14. 144 sections of casing
- 15. 28 in.
- 16. Yes; approximately 8 cm
- 17. 7 homes
- 18. a) Approximately 65 hectares
b) Approximately 259 hectares

Chapter 1: Checkpoint 1, page 25

- 3. a) 26 yd. 2 ft. b) 5280 yd.
c) 84 in.
- 4. Sidney
- 7. Answers will vary depending on the conversion ratios used.
- a) 14 yd. 1 ft. b) 122 cm
c) 1 mi. 427 yd. d) 273 yd. 1 ft. 3 in.
e) 330.2 m f) 5 ft. 9 in.
- 8. 10 ft. of laminate

1.4 Surface Areas of Right Pyramids and Right Cones, page 34

- 4. a) 132 in.² b) 220 cm²
- 5. a) 168 in.² b) 294 cm²
- 6. a) 101 in.² b) 1649 cm²
- 7. a) 151 in.² b) 2356 cm²
- 8. a) 896 cm² b) 628 yd.²
- 9. a)




- b) 7008 ft.²


10. $923\,285\text{ ft.}^2$
 11. a) 2261.9 cm^2
 b) \$11.94
 12. 1520 cm^2
 13. a) 87 m^2
 b) 176 ft.^2
 14. 2.0 m^2 ; I assumed the hides had equal areas.
 15. 188 ft.^2
 16. a) 69.0 mm
 b) 7.6 m
 17. a) Right square pyramid and right cone
 b) Right rectangular prism
 18. The Louvre
 19. a) 193.7 cm^2
 b) 34.9 m^2
 20. 61 ft.^2
 21. 16.0 cm

1.5 Volumes of Right Pyramids and Right Cones, page 42

4. a) 288 yd.^3
 b) 1920 ft.^3
 5. a) 96 yd.^3




- b) 640 ft.^3




6. a) 1571 cm^3
 b) 804 m^3


7. a) 524 cm^3



- b) 268 m^3




8. a) 18 m^3 b) 168 yd.^3
 9. a) 37.7 m^3 b) 2948.9 cm^3
 10. a)




- b) 231.2 m^3

11. a)



- b) 50 yd.^3
 12. 0.3 m^3
 13. b) 441.2 cm^3
 14. a) 5 in.^3
 b) \$3.33
 c) Approximately 7 in.^3

15. a)



- b) 3.8 m c) 15.3 m³
16. 401 ft.³
17. a) 15 cm² b) 23 cm³
c) No, there is also some air inside the tea bag.
18. a) 4.7 cm b) 10.5 m
c) 3.3 m d) 7.4 cm
19. b) 8.0 cm
20. a) 22.9 kL b) Approximately 8.3 kL
21. 10 yd.
22. 49.6 m³

1.6 Surface Area and Volume of a Sphere, page 51

3. a) 314 cm² b) 32 m²
c) 201 ft.² d) 99 cm²
4. a) 524 cm³ b) 17 m³
c) 268 ft.³ d) 92 cm³
5. a) 339 m², 452 m³ b) 191 yd.², 191 yd.³
7. 886.7 m, 2482.7 m³
8. 3.2 cm
9. 12 in.
10. a) 2.1 L b) 8 cups
11. a) Hemisphere b) Hemisphere
12. a) 784 m² b) 2065 kL
13. a) 511 185 933 km²
b) 357 830 153 km²
c) 1 086 781 293 000 km³
d) 1 078 037 876 000 km³
14. Approximately 1 082 696 932 000 km³;
approximately 1 093 440 264 000 km³
15. 239 spheres
16. a) 11 cm; 5 in. b) 1387 cm²; 277 in.²
c) 4855 cm³; 434 in.³ d) Basketball
17. a) 16.4 m³ b) 1.0 m²
18. 529.6 m²; 882.2 m³
19. 42 pumps
20. 45 cookies


21. a) Approximately 69%
b) Assumptions: Ball is created from one solid piece and has greatest possible diameter.
22. $SA = \pi d^2$; $V = \frac{1}{6} \pi d^3$
23. Approximately 5 in.
24. a) Inflated balloon's circumference is 3 times as great
b) Inflated balloon's surface area is 9 times as great
c) Inflated balloon's volume is 27 times as great

Chapter 1: Checkpoint 2, page 54

1. a) 80 ft.² b) 21 m²
c) 1127 m²
2. 425 m²
3. 183 in.²
4. a) 41 ft.³ b) 6 m³
c) 1947 m³
5. a) 9.5 cm b) 2.7 m
c) 17.4 cm
6. a) 973.1 km², 2854.5 km³
b) 109.0 cm², 82.3 cm³
7. 7946 cm²

1.7 Solving Problems Involving Objects, page 59

3. a) 170 cm² b) 1040 ft.²
c) 95 in.² d) 314 in.²
4. a) Object in part c b) Approximately 38 in.³
5. a) 273.3 cm², 353.4 cm³ b) 12.0 m², 2.5 m³
6. a) $5\frac{4}{5}$ in. b) 6.7 cm
7. a)




- b) 2413 cm² c) 6612 cm³
- d) Approximately 2204 cm³, or 2204 mL

8. 93 cm^3
 9. a) Circular-based bin
 b) Square-based bin
 10. a) 1300.0 cm^3 b) 6.2 m^3
 11. a) 856.2 cm^2 b) 24.2 m^2
 12. Approximately 26.4 m^2
 13. a) 1060 in.^3 b) 15 in. by 15 in. by 12 in.
 c) 1820 in.^3


Chapter 1: Review, page 64

1. Answers may vary. For example:
 a) Inch b) Foot
 c) Yard
 3. a) 42 ft. b) 8800 yd.
 c) 75 in. d) 3 yd. 1 ft. 3 in.
 4. 320 in., or 8 yd. 2 ft. 8 in.
 6. Answers will vary depending on the conversion ratios used.
 a) 8 ft. 7 in. b) 136 yd. 2 ft. 1 in.
 c) 3 mi. 1282 yd. d) 1 ft. 2 in.
 7. Answers will vary depending on the conversion ratios used.
 a) 12.5 m b) 6.8 km
 c) 48.3 cm d) 215.9 mm
 8. Answers will vary depending on the conversion ratio used.
 670 750 strides
 9. a) 75 ft.^2 b) 85 cm^2
 c) 898 mm^2 d) 192 m^2
 10. 160 yd.^2
 11. a)



- b) 2.0 m
 c) 6 m^2
 12. a) $8\frac{7}{10} \text{ in.}$ b) 173 in.^2
 13. 125.8 cm^2
 14. 5810 ft.^2
 15. a) 11 m^3 b) 8822 in.^3
 c) 7 ft.^3 d) 221 mm^3

16. No; approximately 132.7 cm^3
 17. 12 cm
 18. a) 24 in.^3 b) 6 in.
 19. a) 2.1 m b) 2.3 cm
 20. a) 254 in.^2 , 382 in.^3
 b) 133 m^2 , 144 m^3
 21.



- a) 763 ft.^2 b) 1527 ft.^3
 22. $4\frac{3}{5} \text{ in.}$
 23. Approximately 98 cm^3
 24. 523 in.^3
 25. a) 480 cm^2 , 595 cm^3 b) 108 ft.^2 , 84 ft.^3
 26. a) $113\,981 \text{ cm}^3$ b) $11\,878 \text{ cm}^2$
 27. a) 8 cm b) 10 mm

Chapter 1: Practice Test, page 67

1. B
 2. C
 3. The volume of the right cylinder is 3 times the volume of the right cone.
 4. a) 28.3 cm^3 , 69.3 cm^2
 b) 1215.8 m^3 , 647.2 m^2
 5. a) A ruler with inches marked
 6. 5.8 cm


Chapter 2 Trigonometry, page 68

2.1 The Tangent Ratio, page 75


3. a) $\tan A = \frac{6}{7}$; $\tan C = \frac{7}{6}$
 b) $\tan D = \frac{3}{2}$; $\tan F = \frac{2}{3}$
 c) $\tan H = \frac{5}{4}$; $\tan J = \frac{4}{5}$
 d) $\tan K = \frac{5}{7}$; $\tan M = \frac{7}{5}$
 4. a) 14° b) 51°
 c) 68° d) 87°
 5. a) 27° b) 45°
 c) 61° d) 69°

6. Sketches will vary. For example:


a)




b)




c)




d)



e)



f)



7. a) $\tan 60^\circ > 1$ b) $\tan 30^\circ < 1$

8. a) 36.4° b) 68.0°

9. b) i) $\angle A \doteq 26.6^\circ$; $\angle B \doteq 63.4^\circ$

ii) $\angle D \doteq 63.4^\circ$; $\angle F \doteq 26.6^\circ$

iii) $\angle G \doteq 63.4^\circ$; $\angle H \doteq 26.6^\circ$

c) No

10. a) 36.0° b) 49.1°

c) 20.3° d) 82.4°

11. a) 11° b) 14°

c) 6° d) 9°

12. Whitehorse

13. $\angle P = \angle RQS \doteq 67.4^\circ$, $\angle R = \angle PQS \doteq 22.6^\circ$

14. 22°

15. 20.6° ; 69.4°

16. The side opposite the acute angle has the same length as the side adjacent to the angle.

17. 25°

18. 22°

19. 146°

20. 76°

21. $\angle X \doteq 50.1^\circ$, $\angle Y = \angle Z \doteq 64.9^\circ$

22. a) There is no least possible value; the tangent can be arbitrarily close to zero.

b) There is no greatest possible value; the tangent can be arbitrarily large.

23. a) 1 ; $\frac{1}{\sqrt{2}}$; $\frac{1}{\sqrt{3}}$; $\frac{1}{\sqrt{4}}$, or $\frac{1}{2}$; $\frac{1}{\sqrt{5}}$

b) $\frac{1}{\sqrt{100}}$, or $\frac{1}{10}$

2.2 Using the Tangent Ratio to Calculate Lengths, page 82

3. a) 2.5 cm

b) 1.4 cm

c) 5.0 cm

d) 7.5 cm

4. a) 2.2 cm

b) 2.8 cm

c) 2.8 cm

5. a) 5.6 cm

b) 4.1 cm

c) 3.8 cm

6. 22.8 m

7. 3.8 m


8. 187 m

9. a) 3.6 cm

b) 10.0 cm

10. Approximately 30 m

11. a)



b) 3.4 cm

12. 40.3 cm^2

13. Approximately 60 m

14. Approximately 58 m, assuming the balloon is directly over the store

15. $\angle QRT = \angle SRT = 26.5^\circ$, $\angle QRS = 53.0^\circ$,
 $\angle QPT = \angle SPT = 56.3^\circ$, $\angle QPS = 112.6^\circ$,
 $\angle RQT = \angle RST = 63.5^\circ$,
 $\angle PQT = \angle PST = 33.7^\circ$,
 $\angle PQR = \angle PSR = 97.2^\circ$,
 $\angle PTQ = \angle PTS = \angle QTR = \angle RTS = 90.0^\circ$
 $PQ = PS \doteq 3.6 \text{ cm}$, $QR = SR \doteq 6.7 \text{ cm}$

16. a) Approximately 38.7°

b) Approximately 63.4°

2.3 Math Lab: Measuring an Inaccessible Height, page 86



- The sum of the angle shown on the protractor and the angle of inclination is 90° .
- 13.5 m
- 25 m

Chapter 2: Checkpoint 1, page 88



- a) 14° b) 56°
c) 53°
- a) 11.2 cm b) 7.3 cm
c) 11.7 cm
- Approximately 23.7 m

2.4 The Sine and Cosine Ratios, page 95



- a) i) Opposite: GH; adjacent: AG; hypotenuse: AH
ii) Opposite: TK; adjacent: AK; hypotenuse: AT
b) i) $\sin A = 0.60$; $\cos A = 0.80$
ii) $\sin A = 0.28$; $\cos A = 0.96$
- a) $\sin 57^\circ \doteq 0.84$; $\cos 57^\circ \doteq 0.54$
b) $\sin 5^\circ \doteq 0.09$; $\cos 5^\circ \doteq 1.00$
c) $\sin 19^\circ \doteq 0.33$; $\cos 19^\circ \doteq 0.95$
d) $\sin 81^\circ \doteq 0.99$; $\cos 81^\circ \doteq 0.16$
- a) 14° b) 50°
c) 33° d) 39°
- a) 34° b) 35°
c) 39° d) 33°
- a) 41° b) 78°
c) 26° d) 66°
- Sketches will vary. For example:




b)



c)



d)



- a) $\angle C \doteq 16.3^\circ$, $\angle D \doteq 73.7^\circ$
b) $\angle F \doteq 63.9^\circ$, $\angle H \doteq 26.1^\circ$
c) $\angle J \doteq 38.0^\circ$, $\angle K \doteq 52.0^\circ$
d) $\angle P \doteq 49.3^\circ$, $\angle Q \doteq 40.7^\circ$

11. 1.3°


12. 79.4°

13. 61°

14. 31°

- a) i) 0.1736... ii) 0.3420...
iii) 0.6427... iv) 0.7660...
v) 0.8660... vi) 0.9848...

16.



The opposite and adjacent sides of an acute angle have the same length, so $\frac{\text{opposite}}{\text{hypotenuse}} = \frac{\text{adjacent}}{\text{hypotenuse}}$.

17. 40°

- a) i) 1 ii) 0
iii) 0 iv) 1

2.5 Using the Sine and Cosine Ratios to Calculate Lengths, page 101

- a) 3.1 cm b) 1.5 cm
c) 1.5 cm d) 3.7 cm
- a) 1.7 cm b) 3.2 cm
c) 5.4 cm d) 7.9 cm
- a) 25.3 cm b) 8.0 cm
c) 7.7 cm d) 12.4 cm

6. 29.7 m
 7. a) 48.3 m
 b) The surveyor could use the tangent ratio or the Pythagorean Theorem.
 8. 4.0 km
 9. 2813 m
 10. 18.3 cm by 4.6 cm
 11. a) 423 cm b) 272 cm
 12. a) i) 21.0 cm ii) 15.1 cm
 13. 186 mm
 14. a) Approximately 139 ft.
 b) $17\,407\text{ ft.}^2$

Chapter 2: Checkpoint 2, page 104


1. a) 30° b) 48°
 c) 56°
 2. 13°
 3. a) i) 0.9848... ii) 0.9396...
 iii) 0.8660... iv) 0.7660...
 v) 0.6427... vi) 0.5
 vii) 0.3420... viii) 0.1736...
 4. a) 4.2 cm b) 2.7 cm
 c) 14.0 cm
 5. Approximately 3.2 km

2.6 Applying the Trigonometric Ratios, page 111


3. a) Sine b) Tangent
 c) Cosine d) Tangent
 4. a) 4.6 cm; cosine b) 4.7 cm; tangent
 c) 11.8 cm; sine d) 14.5 cm; cosine
 5. a) Pythagorean Theorem b) Sine ratio
 c) Pythagorean Theorem d) Pythagorean Theorem
 6. a) $\angle T = 57^\circ$, $TU \doteq 23.0\text{ cm}$, $VU \doteq 19.2\text{ cm}$
 b) $\angle Y = 43^\circ$, $WY \doteq 8.7\text{ cm}$, $XY \doteq 6.3\text{ cm}$
 c) $ZB \doteq 11.3\text{ cm}$, $\angle B \doteq 60.3^\circ$, $\angle Z \doteq 29.7^\circ$
 d) $\angle E = 61^\circ$, $CD \doteq 12.0\text{ cm}$, $CE \doteq 6.6\text{ cm}$
 7. a) 1147 cm b) 1144 cm
 8. 173 ft.
 9. a) 68 km b) 31°
 10. a) 4° b) 15.0 m
 11. a) 31° b) 118°
 12. a) 13.5 cm; 7.8 cm^2 b) 28.9 cm; 47.5 cm^2
 13. 7.3 cm
 14. a) 3 in.^2 b) 15 in.^3
 15. 36 cm
 16. 15.6 cm; 11.6 cm^2

2.7 Solving Problems Involving More than One Right Triangle, page 118

3. a) 6.0 cm b) 6.0 cm
 c) 4.3 cm d) 3.6 cm
 4. a) 5.7 cm b) 4.9 cm
 c) 5.7 cm
 5. a) 93.2° b) 123.7°
 c) 11.1° d) 15.0°
 6. 15 m, 19 m
 7. 51° , 65° , 65°
 8. a) 19 ft. b) 21 ft.
 9. 35 m, 58 m
 10. Approximately 126° , approximately 54°
 11. 4.5 m
 12. a) 53 m b) 29 m
 c) 50 m
 13. a) 5.0 m b) 51.3°
 c) 2.4 m
 14. a) 23 m b) 20 m
 16. a)




- b) 5.1 m
 17. a) 98.1° , 51.7° , 105.1° , 105.1°
 b) 100 mm
 18. a)




- b) 24.0 m
 19. a) 5.4 cm b) 33.9°
 20. Approximately 8.3 m
 21. Approximately 18 in.

Chapter 2: Review, page 124

1. a) 35° b) 65°
2. a) $\tan 20^\circ < 1$ b) $\tan 70^\circ > 1$
3. 6°
4. The triangle is an isosceles right triangle.



5. a) i) 3.7 cm
ii) 3.0 cm
b) Could also use trigonometric ratios
i) Approximately 4.2 cm
ii) Approximately 4.0 cm
6. 327 m
7. a) 11.7 cm b) 13.0 cm
8. 17.5 m
9. 30 m
11. a) 73° ; cosine b) 28° ; sine
- 12.



- a) i) $\frac{5}{13}$ ii) $\frac{12}{13}$
iii) $\frac{5}{13}$ iv) $\frac{12}{13}$
- b) $\sin D = \cos B$; $\sin B = \cos D$
13. 64.2°
14. 36.9°
15. a) 3.9 cm; cosine
b) 4.4 cm; sine
c) 4.7 cm; sine
d) 4.5 cm; cosine
16. 6.0 km
17. 1.6 cm by 2.8 cm
18. a) $CE \doteq 5.0$ cm, $\angle E \doteq 57.3^\circ$, $\angle C \doteq 32.7^\circ$
b) $\angle H = 52^\circ$, $GH \doteq 2.7$ cm, $FH \doteq 4.3$ cm
c) $\angle K = 63^\circ$, $JM \doteq 3.9$ cm, $KM \doteq 2.0$ cm
19. 85.9°
20. a) 35.5 cm; 52.1 cm²
b) 13.0 cm; 10.2 cm²

21. a) 3.2 m b) 8.2 m
22. a) 13.6 cm b) 11.3 cm
c) 21.0°
23. 2316 ft.

Chapter 2: Practice Test, page 127

1. B
2. C
4. $\angle D = 27.0^\circ$, $DE \doteq 6.9$ cm, $EF \doteq 3.5$ cm
5. 203 cm
6. 75.5 m

Cumulative Review Chapters 1 and 2, page 130

1. a) 23 yd. 1 ft. b) \$59.76
2. 276 km
4. Answers will vary depending on the conversion ratios used.
a) 823 cm b) 279 400 m
c) 3 mi. d) 5 ft. 3 in.
5. Answers will vary depending on the conversion ratio used.
The road above The Narrows is higher by approximately 5 ft., or 1.5 m.
6. a) 342 m² b) 208 ft.²
7. 192 ft.³
8. Approximately 6 yd.
9. No
10. a) Hemisphere; 138 in.²
b) Sphere; 3824 in.³
11. 191 m², 170 m³
12. 4478 in.²
13. 222.1 mm², 239.6 mm³
14. a) 31.0° b) 62.5°
15. 26 yd.
16. 201 ft.
17. a) 61.9° b) 68.4°
18. 22°
19. 50 ft. by 94 ft.
20. a) $\angle S = 24.0^\circ$, $RT \doteq 6.4$ m, $RS \doteq 14.4$ m
b) $\angle M = 46.0^\circ$, $MN \doteq 7.1$ cm, $MP \doteq 10.3$ cm
21. 59°
22. $x = 20.0$ cm; $y \doteq 40.0$ cm;
 $\angle PRQ = 46.4^\circ$; $\angle PRS = 133.6^\circ$;
 $\angle PSR = 31.7^\circ$; $\angle QPR = 43.6^\circ$;
 $\angle QPS = 58.3^\circ$; $\angle QRS = 180.0^\circ$;
 $\angle RPS = 14.7^\circ$

Chapter 3 Factors and Products, page 132

3.1 Factors and Multiples of Whole Numbers, page 140

3. a) 6, 12, 18, 24, 30, 36
b) 13, 26, 39, 52, 65, 78
c) 22, 44, 66, 88, 110, 132
d) 31, 62, 93, 124, 155, 186
e) 45, 90, 135, 180, 225, 270
f) 27, 54, 81, 108, 135, 162
4. a) 2, 5 b) 3, 5
c) 3 d) 2, 3, 5
e) 2, 5, 7 f) 2, 3
5. a) $3 \cdot 3 \cdot 5$, or $3^2 \cdot 5$
b) $2 \cdot 2 \cdot 2 \cdot 2 \cdot 5$, or $2^4 \cdot 5$
c) $2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 3$, or $2^5 \cdot 3$
d) $2 \cdot 61$
e) $2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 5$, or $2^5 \cdot 5$
f) $3 \cdot 5 \cdot 13$
6. a) $2^3 \cdot 3 \cdot 5^2$ b) $2 \cdot 5^2 \cdot 23$
c) $2 \cdot 7 \cdot 73$ d) $2 \cdot 3^2 \cdot 5^3$
e) $2^2 \cdot 3^2 \cdot 5^3$ f) $5^3 \cdot 7^2$
8. a) 2 b) 2^3 , or 8
c) 3^3 , or 27 d) 2^2 , or 4
e) 2^5 , or 32 f) $2^2 \cdot 5$, or 20
9. a) 5 b) $2^3 \cdot 5$, or 40
c) $2 \cdot 3 \cdot 7$, or 42 d) 2^2 , or 4
10. a) $2^2 \cdot 3 \cdot 7$, or 84 b) $3^2 \cdot 5 \cdot 7$, or 315
c) $2^2 \cdot 3^2 \cdot 5$, or 180 d) $2 \cdot 3 \cdot 7 \cdot 19$, or 798
e) $2^5 \cdot 3^2 \cdot 5$, or 1440 f) $2^2 \cdot 7 \cdot 13$, or 364
11. a) $2^2 \cdot 3^2 \cdot 5 \cdot 19$, or 3420
b) $2^5 \cdot 3 \cdot 5 \cdot 11$, or 5280
c) $2^2 \cdot 3^2 \cdot 5^2$, or 900
d) $2^3 \cdot 3^3 \cdot 5$, or 1080
12. Greatest common factor: 2;
least common multiple: $2^2 \cdot 3 \cdot 7$, or 84
13. 2 · 3, or 6
14. The greatest common factor of the two numbers is 1.
15. a) $\frac{37}{65}$ b) $\frac{17}{19}$
c) $\frac{13}{18}$ d) $\frac{42}{61}$
e) $\frac{49}{110}$ f) $\frac{33}{17}$

16. a) $\frac{149}{112}$ b) $\frac{65}{60}$, or $\frac{13}{12}$
c) $\frac{43}{264}$ d) $\frac{304}{210}$, or $\frac{152}{105}$
e) $\frac{121}{600}$ f) $\frac{239}{90}$
g) $\frac{27}{20}$ h) $\frac{77}{12}$

17. 800 m
18. No; 1 does not have any prime factors.
19. a) 72 cm by 72 cm b) Yes
20. a) Yes b) Yes
c) 660 feet
21. Yes
22. 30 cm

3.2 Perfect Squares, Perfect Cubes, and Their Roots, page 146

4. a) 14 b) 16
c) 19 d) 17
e) 21
5. a) 7 b) 8
c) 10 d) 11
e) 15
6. a) Perfect square
b) Perfect square and perfect cube
c) Neither
d) Perfect square
e) Perfect square and perfect cube
f) Perfect cube
7. a) 22 mm b) 42 yd.
8. a) 18 in. b) 25 ft.
9. 96 ft.²
10. 35 937 ft.³
11. No; 2000 is not a perfect cube.
12. These answers assume that the endpoints of each range are included in the range.
a) Perfect squares: 324, 361; perfect cube: 343
b) Perfect squares: 676, 729; perfect cube: 729
c) Perfect squares: 841, 900
d) Perfect squares: 1225, 1296; perfect cube: 1331
13. The first 5 are: 0, 1, 64, 729, 4096
14. 12 ft.
15. a) $\frac{45x^2}{8}$ b) $x = 4$

16. Edge length: 6 units

17. a) $11x^2y$

b) $4x^2y$

18. $1^3 + 12^3, 9^3 + 10^3$

Chapter 3: Checkpoint 1, page 149

1. a) $2^2 \cdot 3^2 \cdot 5 \cdot 7$

b) $2^7 \cdot 3 \cdot 11$

c) $2^3 \cdot 3^2 \cdot 5 \cdot 17$

d) $5 \cdot 11 \cdot 19$

e) $2^4 \cdot 3^3 \cdot 7$

f) $3 \cdot 5^2 \cdot 7^2$

2. a) 2^3 , or 8

b) $2^2 \cdot 3$, or 12

c) 5

d) 2^4 , or 16

e) 2^3 , or 8

f) 5^2 , or 25

3. a) $2^2 \cdot 3 \cdot 5 \cdot 7$, or 420

b) $2^5 \cdot 3 \cdot 5$, or 480

c) $2^3 \cdot 3^2 \cdot 5$, or 360

d) $2^5 \cdot 3 \cdot 5$, or 480

e) $2^6 \cdot 7^2$, or 3136

f) $2 \cdot 3 \cdot 5^2 \cdot 11$, or 1650

4. a) $\frac{103}{33}$

b) $\frac{71}{35}$

c) $\frac{27}{70}$

5. 18 980 days; 52 years

6. a) 20

b) 28

c) 24

d) 33

e) 39

f) 55

7. a) 12

b) 15

c) 20

d) 18

e) 22

f) 21

8. a) Neither

b) Perfect square

c) Perfect square and perfect cube

d) Perfect square

e) Perfect cube

f) Neither

9. a) Perfect squares: 400, 441, 484

b) Perfect squares: 900, 961; perfect cube: 1000

c) Perfect square: 1156

10. 26 cans

3.3 Common Factors of a Polynomial, page 155

Gray algebra tiles represent positive tiles and black tiles represent negative algebra tiles.

4. a) $3x + 12$; 3, $x + 4$

b) $4x^2 + 10x$; $2x$, $2x + 5$

c) $12x^2 - 8x + 16$; 4, $3x^2 - 2x + 4$

5. a) 3

b) m

6. a) i) $3(2 + 5n)$

ii) $3(2 - 5n)$

iii) $3(5n - 2)$

iv) $3(-5n + 2)$


b) i) $m(4 + m)$

ii) $m(m + 4)$


iii) $m(4 - m)$

iv) $m(m - 4)$

7. a) $5(y + 2)$




b) $6(1 + 2x^2)$



c) $3(3k + 2)$



d) $2s(2s + 7)$



e) $y(1 + y)$



f) $h(3 + 7h)$



8. a) $3b^2(3 - 4b)$

b) $12(4s^3 - 1)$


c) $-a^2(1 + a)$

d) $3x^2(1 + 2x^2)$

e) $4y(2y^2 - 3)$

f) $-7d(1 + 2d^3)$


9. a) $3(x^2 + 4x - 2)$




b) $2(2 - 3y - 4y^2)$



c) $-7(m + m^2 + 2)$




d) $2(5n - 3 - 6n^2)$



e) $2(4 + 5x + 3x^2)$



f) $-3(3 - 4b - 2b^2)$



10. a) $5(1 + 3m^2 - 2m^3)$

c) $v(6v^3 + 7 - 8v^2)$

e) $6x(4 + 5x - 2x^3)$

b) $9(3n + 4 - 2n^3)$

d) $-c^2(3 + 13c^2 + 12c)$

f) $s(s^3 + s - 4)$

11. a) $-12x^2 + 20x$

b) $4x$ and $(-3x + 5)$

c) The factors are the dimensions of the rectangle.

12. a) i) $3m(m + 3m^2 - 1)$

ii) $-4(4 - 2n + n^3)$

b) Expanded his solutions

13. The monomial is 1 when the term is the common factor.
The monomial is -1 when the term has the opposite sign of the common factor.

14. a) $4x - 4 = 4(x - 1)$

b) $16m^2 - 24m - 16 = 8(2m^2 - 3m - 2)$

c) $-8n^3 - 6n^2 - 10n = -2n(4n^2 + 3n + 5)$

15. a) i) $2 \cdot 2 \cdot s \cdot t \cdot t$, or $4st^2$

ii) $a \cdot a \cdot b$, or a^2b

iii) $2 \cdot 2 \cdot 3 \cdot x \cdot x \cdot y \cdot y$, or $12x^2y^2$

b) i) $4st^2(s + 3st + 9)$

ii) $4st^2(3st - s - 9)$

iii) $-a^2b(3a + 9a^2 - 8)$

iv) $a^2b(9a^2 + 3a - 8)$

v) $12x^2y^2(3y^2 + x + x^2y)$

vi) $-12x^2y^2(3y^2 + x^2y + x)$

16. a) $5x(5y + 3x - 6xy^2)$

b) $3mn(17m + 13n - 24)$

c) $3p^2q^2(3p^2 - 2pq + 4q^2)$

d) $a^2b^2(10a + 12b^2 - 5)$

e) $4cd(3d - 2 - 5c)$

f) $7rs^2(r^2s + 2r - 3)$

17. a) $SA = 2\pi r(r + h)$

b) Approximately 2639 cm^2

18. a) $SA = \pi r(r + s)$

b) Approximately 679 cm^2

19. a) Assume the area of the base of the silo is not included in the surface area. $SA = 2\pi rh + 2\pi r^2$;

$SA = 2\pi r(h + r)$; approximately 603 m^2

b) $V = \pi r^2 h + \frac{2}{3}\pi r^3$; $V = \pi r^2 \left(h + \frac{2}{3}r \right)$;

approximately 1583 m^3

20. Yes

21. a) $\frac{2\pi rh}{2\pi r^2 + 2\pi rh}$

b) $\frac{h}{r + h}$

22. a) 2; 3

b) $n - 3$

c) $\frac{n^2}{2} - \frac{3n}{2} = \frac{n}{2}(n - 3)$

3.4 Math Lab: Modelling Trinomials as Binomial Products, page 158

1. a) Can be represented




- b) Can be represented



- c) Cannot be represented
d) Cannot be represented
e) Cannot be represented
f) Cannot be represented


2. a) Can be represented



- b) Can be represented



- c) Cannot be represented
d) Cannot be represented
e) Cannot be represented
f) Can be represented



3. 7, 8, 13
4. 4, 7, 9, 10


3.5 Polynomials of the Form $x^2 + bx + c$, page 166

4. a) $(x+1)(x+3) = x^2 + 4x + 3$
b) $(x+2)(x+4) = x^2 + 6x + 8$
c) $(x+5)(x+5) = x^2 + 10x + 25$
d) $(x+3)(x+6) = x^2 + 9x + 18$


5. a) $b^2 + 7b + 10$



- b) $n^2 + 11n + 28$



- c) $h^2 + 11h + 24$



- d) $k^2 + 7k + 6$



6. a) i) $x^2 + 4x + 4$

ii)



- iii) $(x+2)(x+2)$

- b) i) $x^2 + 5x + 4$


ii)



- iii) $(x+1)(x+4)$

- c) i) $x^2 + 6x + 8$

ii)



- iii) $(x+2)(x+4)$

- d) i) $x^2 + 7x + 12$

ii)



- iii) $(x+3)(x+4)$

7. a) i) 1, 2 ii) 2, 3
 iii) 1, 9 iv) 2, 5
 v) 3, 4 vi) 3, 5
 b) i) $(v+1)(v+2)$ ii) $(w+2)(w+3)$
 iii) $(s+1)(s+9)$ iv) $(t+2)(t+5)$
 v) $(y+3)(y+4)$ vi) $(h+3)(h+5)$


8. a) i) $(v+1)(v+1)$




- ii) $(v+2)(v+2)$



- iii) $(v+3)(v+3)$



- iv) $(v+4)(v+4)$



- b) The rectangles are squares; the binomial factors are the same.

c) $v^2 + 10v + 25 = (v+5)(v+5)$;

$v^2 + 12v + 36 = (v+6)(v+6)$;

$v^2 + 14v + 49 = (v+7)(v+7)$

9. Area models and/or rectangle diagrams may vary.

For example:

- a) $m^2 + 13m + 40$

	m	8
m	$(m)(m) = m^2$	$(m)(8) = 8m$
5	$(5)(m) = 5m$	$(5)(8) = 40$

- b) $y^2 + 12y + 27$

	y	3
y	$(y)(y) = y^2$	$(y)(3) = 3y$
9	$(9)(y) = 9y$	$(9)(3) = 27$

- c) $w^2 + 18w + 32$

	w	16
w	$(w)(w) = w^2$	$(w)(16) = 16w$
2	$(2)(w) = 2w$	$(2)(16) = 32$

- d) $k^2 + 14k + 13$

	k	1
k	$(k)(k) = k^2$	$(k)(1) = k$
13	$(13)(k) = 13k$	$(13)(1) = 13$

10. a) $(w+3)(w+2) = w^2 + 5w + 6$

b) $(x+5)(x+2) = x^2 + 7x + 10$

c) $(y+10)(y+2) = y^2 + 12y + 20$

11. a) $(x+4)(x+6)$ b) $(m+2)(m+8)$

c) $(p+1)(p+12)$ d) $(s+2)(s+10)$

e) $(n+1)(n+11)$ f) $(h+2)(h+6)$

g) $(q+1)(q+6)$ h) $(b+2)(b+9)$

12. a) $g^2 + 4g - 21$

	g	7
g	$(g)(g) = g^2$	$(g)(7) = 7g$
-3	$(-3)(g) = -3g$	$(-3)(7) = -21$

b) $h^2 - 5h - 14$

	h	-7
h	$(h)(h) = h^2$	$(h)(-7) = -7h$
2	$(2)(h) = 2h$	$(2)(-7) = -14$

c) $22 - 13j + j^2$

	2	$-j$
11	$(11)(2) = 22$	$(11)(-j) = -11j$
$-j$	$(-j)(2) = -2j$	$(-j)(-j) = j^2$

d) $k^2 + 8k - 33$

	k	11
k	$(k)(k) = k^2$	$(k)(11) = 11k$
-3	$(-3)(k) = -3k$	$(-3)(11) = -33$

e) $84 - 5h - h^2$

	7	$-h$
12	$(12)(7) = 84$	$(12)(-h) = -12h$
h	$(h)(7) = 7h$	$(h)(-h) = -h^2$

f) $m^2 - 81$

	m	9
m	$(m)(m) = m^2$	$(m)(9) = 9m$
-9	$(-9)(m) = -9m$	$(-9)(9) = -81$

g) $n^2 - 18n + 56$

	n	-4
n	$(n)(n) = n^2$	$(n)(-4) = -4n$
-14	$(-14)(n) = -14n$	$(-14)(-4) = 56$

h) $p^2 - 11p - 102$

	p	-17
p	$(p)(p) = p^2$	$(p)(-17) = -17p$
6	$(6)(p) = 6p$	$(6)(-17) = -102$

13. a) $r^2 - 9r - 52$

b) $s^2 - 20s + 75$

14. a) $(b-1)(b+20)$

b) $(t-3)(t+18)$

c) $(x-2)(x+14)$

d) $(n+3)(n-8)$

e) $(a+4)(a-5)$

f) $(y+6)(y-8)$

g) $(m-5)(m-10)$

h) $(a-6)(a-6)$

15. a) $(1+k)(12+k)$

b) $(2+g)(-8+g)$

c) $(5+y)(12+y)$

d) $(9+z)(8-z)$

16. a) i) $x^2 + 3x + 2$; 132

ii) $x^2 + 4x + 3$; 143

b) The coefficients of the terms of the polynomial are the digits in the product of integers.

17. a) $(m+5)(m-12)$

b) $(w-5)(w-9)$

c) $(b-3)(b+12)$

18. a) i) $t^2 + 11t + 28$

ii) $t^2 - 11t + 28$

iii) $t^2 + 3t - 28$

iv) $t^2 - 3t - 28$

b) i) Because the constant terms in the binomials have the same sign

ii) Because the constant terms in the binomials have opposite signs

iii) Add the constant terms in the binomials

19. a) $\pm 7, \pm 11$; 4 integers

b) $0, \pm 8$; 3 integers

c) $\pm 6, \pm 9$; 4 integers

d) $\pm 1, \pm 4, \pm 11$; 6 integers

e) $\pm 9, \pm 11, \pm 19$; 6 integers

f) $0, \pm 6, \pm 15$; 5 integers

20. Infinitely many integers are possible. For example:

- a) 0, -2, -6, -12, -20, -30, ...
- b) 0, -2, -6, -12, -20, -30, ...
- c) 1, 0, -3, -8, -15, -24, -35, ...
- d) 1, 0, -3, -8, -15, -24, -35, ...
- e) 2, 0, -4, -10, -18, -28, -40, ...
- f) 2, 0, -4, -10, -18, -28, -40, ...

- 21. a) $4(y-7)(y+2)$ b) $-3(m+2)(m+4)$
- c) $4(x-3)(x+4)$ d) $10(x+2)(x+6)$
- e) $-5(n-1)(n-7)$ f) $7(c-2)(c-3)$
- 23. a) i) $(h+2)(h-12)$ ii) $(h-2)(h+12)$
- iii) $(h-4)(h-6)$ iv) $(h+4)(h+6)$

b) The first 6 are:

$$h^2 \pm 13h \pm 30, h^2 \pm 15h \pm 54, h^2 \pm 17h \pm 60, \\ h^2 \pm 25h \pm 84, h^2 \pm 20h \pm 96, h^2 \pm 26h \pm 120$$

3.6 Polynomials of the Form $ax^2 + bx + c$, page 177

- 5. a) $(2m+1)(m+3) = 2m^2 + 7m + 3$
- b) $(3p+2)(p+4) = 3p^2 + 14p + 8$
- c) $(3w+1)(2w+1) = 6w^2 + 5w + 1$
- d) $(4v+3)(3v+2) = 12v^2 + 17v + 6$
- 6. a) $2v^2 + 7v + 6$ b) $3r^2 + 13r + 4$
- c) $6g^2 + 13g + 6$ d) $8z^2 + 26z + 15$
- e) $9t^2 + 24t + 16$ f) $4r^2 + 12r + 9$
- 7. a) i) $2x^2 + 5x + 2$


ii)



iii) $(2x+1)(x+2)$

- b) i) $3x^2 + 11x + 6$


ii)



iii) $(x+3)(3x+2)$

- c) i) $3x^2 + 8x + 4$

ii)



iii) $(x+2)(3x+2)$

- d) i) $4x^2 + 9x + 2$

ii)



iii) $(x+2)(4x+1)$

- 8. a) $(2w+1)(w+6) = 2w^2 + 13w + 6$
- b) $(2g-5)(3g-3) = 6g^2 - 21g + 15$
- c) $(-4v-3)(-2v-7) = 8v^2 + 34v + 21$
- 9. a) $15 + 23f + 4f^2$ b) $15 - 29t + 12t^2$
- c) $90 + 11r - 2r^2$ d) $36 - 24m + 4m^2$
- e) $-24 + 50x + 14x^2$ f) $-36 + 60n - 25n^2$
- 10. a) $6c^2 + 23c + 20$ b) $-21t^2 - 32t + 5$
- c) $32r^2 + 48r - 14$ d) $5t^2 + 46t + 9$
- e) $35h^2 + 29h - 30$ f) $-36y^2 + 84y - 49$


- 11. a) i) $(t+1)(3t+1)$




ii) $(t+2)(3t+2)$



iii) $(t+3)(3t+3)$



iv) $(t+4)(3t+4)$



b) The side lengths increase by 1 each time; the constant terms in the binomial factors increase by 1 each time.

- c) $3t^2 + 20t + 25 = (t+5)(3t+5);$
- $3t^2 + 24t + 36 = (t+6)(3t+6);$
- $3t^2 + 28t + 49 = (t+7)(3t+7)$

- 12. a) i) $(n+6)(2n+1)$ ii) $(n-6)(2n-1)$
- b) i) $(n+6)(2n-1)$ ii) $(n-6)(2n+1)$
- c) i) $(n+2)(2n+3)$ ii) $(n-2)(2n-3)$


The trinomials in each pair have middle terms with the same value, but opposite signs. The constant terms in the binomial factors have opposite signs.

13. a) $(v+2)(2v+1)$ b) $(a+4)(2a+3)$
 c) $(k+5)(2k+3)$ d) $(m-4)(2m-3)$
 e) $(k-3)(2k-5)$ f) $(m+7)(2m+1)$
 g) $(g+6)(2g+3)$ h) $(n+6)(2n-3)$
14. a) i) 1, 15 ii) 2, 12
 iii) 3, 5 iv) 3, 4
 v) 1, 12 vi) 3, 8
 b) i) $(v+5)(3v+1)$
 ii) $(m+4)(3m+2)$
 iii) $(b+1)(3b+5)$
 iv) $(a+1)(4a+3)$
 v) $(d+3)(4d+1)$
 vi) $(v+2)(4v+3)$
15. a) $(a-2)(5a+3)$ b) $(v-5)(3v+2)$
 c) $(s+4)(5s-1)$ d) $(2c-3)(7c+1)$
 e) $(2a+5)(4a-1)$ f) $(2r-3)(4r-1)$
 g) $(d+1)(6d-5)$ h) $(3e-2)(5e+1)$
16. a) $(2u+7)(3u-2)$
 b) $(3k-10)(k+3)$
 c) $(4v-5)(v-4)$
17. $(3g+7)(5g-6)$
18. a) $10(r+2)(2r+3)$ b) $5(a-4)(3a-1)$
 c) $3(2h+3)(3h-2)$ d) $6(2u-3)(2u-3)$
 e) $4(m-5)(3m+2)$ f) $2(3g+5)(4g-7)$
19. a) $(2y-1)(7y-3)$ b) $(p-2)(10p+3)$
 c) $(2r-7)(5r+1)$ d) $(3g+1)(5g-2)$
 e) $(2x-3)(2x+5)$ f) $(3d-4)(3d-4)$
 g) $(3t+2)(3t+2)$ h) $(5y+2)(8y-3)$
 i) $(2c+3)(12c-5)$ j) $(2x+5)(4x-3)$
20. These answers do not include cases where there is a common constant factor among the terms of the polynomial.
 a) $\pm 7, \pm 8, \pm 13$; 6 integers
 b) $\pm 20, \pm 25, \pm 29, \pm 52, \pm 101$; 10 integers
 c) $\pm 3, \pm 15, \pm 25, \pm 53$; 8 integers
 d) $\pm 22, \pm 23, \pm 26, \pm 29, \pm 34, \pm 43, \pm 62, \pm 121$; 16 integers
 e) $\pm 6, \pm 10$; 4 integers
 f) ± 1 ; 2 integers
21. a) i) $(r+1)(4r-5)$
 ii) Cannot be factored
 iii) Cannot be factored
 iv) $(w-2)(2w-1)$
 v) $(h-3)(3h+1)$
 vi) Cannot be factored


22. a) i) $(n+2)(3n+5)$ ii) $(n-2)(3n-5)$
 iii) $(n+1)(3n+10)$ iv) $(n-1)(3n-10)$
 v) $(n+5)(3n+2)$ vi) $(n-5)(3n-2)$
 b) Yes; $3n^2+31n+10$ and $3n^2-31n+10$
23. $9m^2 \pm 24m + 16$, $9m^2 \pm 25m + 16$, $9m^2 \pm 26m + 16$,
 $9m^2 \pm 30m + 16$, $9m^2 \pm 40m + 16$, $9m^2 \pm 51m + 16$,
 $9m^2 \pm 74m + 16$, $9m^2 \pm 145m + 16$

Chapter 3: Checkpoint 2, page 180

1. a) $6x+15$; 3 and $(2x+5)$




- b) $4x+12$; 4 and $(x+3)$



2. a) i) $4(a+2)$




- ii) $3(c-2)$




- iii) $-v(2v+5)$



- iv) $2(x^2+7x+3)$




v) $-3(r^2 - 5r + 1)$



vi) $3a(5a^2 - ab - 2b^2)$

vii) $4(3 - 8x + 2x^2)$




viii) $4y(3x^2 - 2x - 4)$

b) The polynomials in part vi and part viii


3. Answers will vary. For example:

$$x^2 + 5x + 6 = (x + 3)(x + 2)$$



4. Answers will vary. For example:

$$3x^2 + 10x + 8 = (x + 2)(3x + 4)$$



5. Area models and rectangle diagrams may vary.

For example:

a) $x^2 + 5x + 4$

	x	4
x	$(x)(x) = x^2$	$(x)(4) = 4x$
1	$(1)(x) = x$	$(1)(4) = 4$

b) $d^2 + d - 6$

	d	3
d	$(d)(d) = d^2$	$(d)(3) = 3d$
-2	$(-2)(d) = -2d$	$(-2)(3) = -6$

c) $x^2 - 6x + 8$

	x	-2
x	$(x)(x) = x^2$	$(x)(-2) = -2x$
-4	$(-4)(x) = -4x$	$(-4)(-2) = 8$

d) $30 - r - r^2$

	6	r
5	$(5)(6) = 30$	$(5)(r) = 5r$
$-r$	$(-r)(6) = -6r$	$(-r)(r) = -r^2$

e) $g^2 + 4g - 5$

	g	-1
g	$(g)(g) = g^2$	$(g)(-1) = -g$
5	$(5)(g) = 5g$	$(5)(-1) = -5$

f) $20 - 12t + t^2$

	10	$-t$
2	$(2)(10) = 20$	$(2)(-t) = -2t$
$-t$	$(-t)(10) = -10t$	$(-t)(-t) = t^2$

6. a) $(s + 5)(s + 6)$

c) $(4 - b)(5 - b)$

e) $(z + 3)(z + 10)$

7. a) $3(x - 2)(x + 7)$

c) $-(3 + m)(8 + m)$

8. a) $2c^2 + 7c + 3$

c) $9f^2 - 9f - 4$

e) $30 - 8r - 6r^2$

9. a) $(j + 4)(2j + 5)$

c) $(k - 4)(5k - 3)$

e) $(2y - 1)(4y + 1)$

b) $(n + 5)(n - 6)$

d) $-(1 + t)(11 - t)$

f) $-(k - 3)(k - 6)$

b) $-2(y - 3)(y - 8)$

d) $(2 - y)(25 + y)$

b) $-4m^2 + 21m - 5$

d) $12z^2 - 20z + 3$

f) $8 + 20h + 8h^2$

b) $(v + 2)(3v - 5)$

d) $(3h + 2)(3h + 4)$

f) $(3 - 4u)(2 - 5u)$

3.7 Multiplying Polynomials, page 186


4. a) $g^3 + 3g^2 + 5g + 3$
 b) $2 + 7t + 6t^2 + 4t^3 + t^4$
 c) $2w^3 + 11w^2 + 26w + 21$
 d) $12 + 29n + 22n^2 + 8n^3 + n^4$
5. a) $6z^2 + 5zy + y^2$
 b) $12f^2 + 4f - 25fg - 3g + 12g^2$
 c) $8a^2 + 22ab + 15b^2$
 d) $12a^2 + 4a - 31ab - 5b + 20b^2$
 e) $4r^2 + 4rs + s^2$
 f) $9t^2 - 12tu + 4u^2$
6. a) i) $4x^2 + 4xy + y^2$
 ii) $25r^2 + 20rs + 4s^2$
 iii) $36c^2 + 60cd + 25d^2$
 iv) $25v^2 + 70vw + 49w^2$
 v) $4x^2 - 4xy + y^2$
 vi) $25r^2 - 20rs + 4s^2$
 vii) $36c^2 - 60cd + 25d^2$
 viii) $25v^2 - 70vw + 49w^2$
 b) i) $p^2 + 6pq + 9q^2$
 ii) $4s^2 - 28st + 49t^2$
 iii) $25g^2 + 40gh + 16h^2$
 iv) $100h^2 - 140hk + 49k^2$
7. a) i) $x^2 - 4y^2$ ii) $9r^2 - 16s^2$
 iii) $25c^2 - 9d^2$ iv) $4v^2 - 49w^2$
 b) i) $121g^2 - 25h^2$ ii) $625m^2 - 49n^2$
8. a) $3y^3 + y^2 - 26y + 16$
 b) $4r^3 - 7r^2 - 14r - 3$
 c) $2b^3 + 17b^2 - 13b + 2$
 d) $3x^3 + 11x^2 - 39x - 7$
9. a) $x^2 + 3x + 2xy + 3y + y^2$
 b) $x^2 + 3x + xy + 2y + 2$
 c) $a^2 + 2ab + b^2 + ac + bc$
 d) $3s + st + 5t + t^2 + 6$
10. a) $x^2 - x - 2y - 4y^2$
 b) $2c^2 + 2c - cd - 3d - 3d^2$
 c) $a^2 - 4a - 3ab + 20b - 10b^2$
 d) $p^2 + 2pq - 8q^2 - pr + 2qr$
11. $2r^2 - 13rs + 12r + 15s^2 - 18s$
12. $x^3 + 10x^2 + 23x + 14$
13. a) $4r^4 + 13r^3 + 12r^2 + 5r + 2$
 b) $2d^4 + 14d^3 + 19d^2 + 12d + 3$
 c) $-4c^4 + 26c^3 - c^2 - 22c - 6$
 d) $8n^4 - 18n^3 - 7n^2 + 16n - 3$
14. $-3g^4 - 7g^3 + 10g^2 + 18g - 8$

15. a) $9s^2 + 41s + 52$
 b) $13x^2 + 4x + 40$
 c) $18m^2 - 2m - 42mn - 4n$
 d) 0
 e) $3x^2 - 28x + 10$
 f) $7a^2 + 2a - 7$
16. a) $20 - 2x$
 b) $10 - 2x$
 c) $4x^2 - 60x + 200$
 d) $4x^3 - 60x^2 + 200x$
17. a) $27x^2 + 43x + 16$
 b) $x^2 + 2x - 2$
18. a) $x^3 - 6x^2 + 12x - 8$
 b) $8y^3 + 60y^2 + 150y + 125$
 c) $64a^3 - 144a^2b + 108ab^2 - 27b^3$
 d) $c^3 + 3c^2d + 3cd^2 + d^3$
19. a) $12a^3 + 2a^2 - 4a$
 b) $-6r^3 + 3r^2 + 3r$
 c) $40x^4 - 50x^3 + 15x^2$
 d) $-8x^3y - 10x^2y + 25xy$
 e) $4b^3 + 2b^2c - 2bc^2$
 f) $y^6 - y^2$
20. a) $(2x + 3)^3 = 8x^3 + 36x^2 + 54x + 27$
 b) $6(2x + 3)^2 = 24x^2 + 72x + 54$
21. a) $6x^3 + 2x^2 - 128x - 160$
 b) $3b^3 - b^2 - 172b + 224$
 c) $18x^3 + 3x^2 - 88x - 80$
 d) $50a^3 - 235a^2 + 228a - 63$
 e) $8k^3 + 12k^2 - 18k - 27$
22. a) $x^3 + 3x^2y + 3xy^2 + y^3 + 3x^2 + 6xy + 3y^2 + 3x + 3y + 1$
 b) $x^3 - 3x^2y + 3xy^2 - y^3 - 3x^2 + 6xy - 3y^2 + 3x - 3y - 1$
 c) $x^3 + 3x^2y + 3xy^2 + y^3 + 3x^2z + 6xyz + 3y^2z + 3xz^2 + 3yz^2 + z^3$
 d) $x^3 - 3x^2y + 3xy^2 - y^3 - 3x^2z + 6xyz - 3y^2z + 3xz^2 - 3yz^2 - z^3$

3.8 Factoring Special Polynomials, page 194

4. a) $x^2 + 4x + 4$ b) $9 - 6y + y^2$
 c) $25 + 10d + d^2$ d) $49 - 14f + f^2$
 e) $x^2 - 4$ f) $9 - y^2$
 g) $25 - d^2$ h) $49 - f^2$
5. a) Difference of squares
 b) Neither
 c) Neither
 d) Perfect square trinomial

6. a) $(x+7)(x-7)$ b) $(b+11)(b-11)$
 c) $(1+q)(1-q)$ d) $(6+c)(6-c)$
7. a) i) $(a+5)^2$ ii) $(b-6)^2$
 iii) $(c+7)^2$ iv) $(d-8)^2$
 v) $(e+9)^2$ vi) $(f-10)^2$
- b) $g^2 + 22g + 121 = (g+11)^2$;
 $h^2 - 24h + 144 = (h-12)^2$;
 $i^2 + 26i + 169 = (i+13)^2$;
 $j^2 - 28j + 196 = (j-14)^2$
8. a) $(2x-3)^2$ b) $(3+5n)^2$
 c) $(9-2v)^2$ d) $(5+4h)^2$
 e) $(3g+8)^2$ f) $(7r-2)^2$
9. a) $x^2; y^2; x^2 - y^2$
 b) $(x-y)$ and $(x+y)$; $(x-y)(x+y)$
10. a) $(3d+4f)(3d-4f)$
 b) $(5s+8t)(5s-8t)$
 c) $(12a+3b)(12a-3b)$, or $9(4a+b)(4a-b)$
 d) $(11m+n)(11m-n)$
 e) $(9k+7m)(9k-7m)$
 f) $(10y+9z)(10y-9z)$
 g) $(v+6t)(v-6t)$
 h) $(2j+15h)(2j-15h)$
11. a) $(y+2z)(y+5z)$ b) $(2w+3x)(2w-7x)$
 c) $(3s-u)(4s-u)$ d) $(t-v)(3t-4v)$
 e) $(2r+3s)(5r-3s)$ f) $(2p+7q)(4p-5q)$
12. Trinomials in parts a, c, and d are perfect squares.
 a) $(2x+7y)^2$ b) $(3m-n)(5m+4n)$
 c) $(4r+t)^2$ d) $(3a-7b)^2$
 e) $(3h+4k)(4h+3k)$ f) $(3f-5g)(5f-2g)$
13. a) $8(m+3n)(m-3n)$
 b) $2(2z+y)^2$
 c) $3(2x+3y)(2x-3y)$
 d) $2(2p+5q)^2$
 e) $-3(2u-v)(4u+3v)$
 f) $-2(3b+8c)(3b-8c)$
14. a)



- b) $\pi R^2 - \pi r^2 = \pi(R+r)(R-r)$
 c) Approximately 314 159 cm^2

15. a) i) ± 14 ii) 25
 iii) 9
 b) i) 2 integers ii) 1 integer
 iii) 1 integer
16. $-2, -1, 0; -1, 0, 1$; 2 possibilities
17. 39 999
18. $5x^2 + 34x + 24$
19. a) i) Neither
 ii) Difference of squares
 iii) Difference of squares
 iv) Perfect square trinomial
- b) ii) $(-10+r)(10+r)$
 iii) $(9ab+1)(9ab-1)$
 iv) $(4s^2+1)^2$
20. a) $(x+2)(x-2)(x+3)(x-3)$
 b) $(a+1)(a-1)(a+4)(a-4)$
 c) $(y+1)(y-1)(y+2)(y-2)$
21. a) $8(d+2e)(d-2e)$
 b) $\frac{1}{4}(10m+n)(10m-n)$, or $\left(5m+\frac{1}{2}n\right)\left(5m-\frac{1}{2}n\right)$
 c) $2y^2(3x+5y)(3x-5y)$
 d) Cannot be factored
 e) Cannot be factored
 f) $\frac{1}{196}(7x+2y)(7x-2y)$, or $\left(\frac{x}{4}+\frac{y}{7}\right)\left(\frac{x}{4}-\frac{y}{7}\right)$

Chapter 3: Review, page 198

1. a) 2, 3, 11; $2 \cdot 3^3 \cdot 11$
 b) 2, 3, 5, 7; $2^2 \cdot 3 \cdot 5^2 \cdot 7$
 c) 3, 5, 13; $3 \cdot 5^3 \cdot 13$
 d) 3, 7, 11, 13; $3^2 \cdot 7 \cdot 11 \cdot 13$
2. a) $2^2 \cdot 5$, or 20 b) $5 \cdot 7$, or 35
 c) 2^4 , or 16 d) 2^2 , or 4
3. a) $2^2 \cdot 3^2 \cdot 5 \cdot 7$, or 1260
 b) $2^3 \cdot 3 \cdot 5 \cdot 13 \cdot 103$, or 160 680
 c) $2^3 \cdot 5^3$, or 1000
 d) $2^4 \cdot 3^2 \cdot 5 \cdot 17$, or 12 240
4. 61 beads
5. a) $\frac{7}{9}$ b) $\frac{11}{17}$
 c) $\frac{13}{15}$ d) $\frac{247}{576}$
 e) $\frac{20}{27}$ f) $\frac{23}{160}$
6. a) 28 in. b) 32 cm

7. a) 12 cm b) 14 ft.

8. a) Perfect square; $\sqrt{256} = 16$

b) Perfect square; $\sqrt{324} = 18$

c) Perfect square and perfect cube;
 $\sqrt{729} = 27$; $\sqrt[3]{729} = 9$

d) Neither

e) Perfect square; $\sqrt{1936} = 44$

f) Perfect cube; $\sqrt[3]{9261} = 21$

9. 540 ft.

10. 44 cm

11. a) $4m(2-m)$ b) $-3(1-3g^2)$

c) $7a^2(4-a)$ d) $3a^2b^2c(2b-5c)$

e) $-6mn(4m+n)$ f) $7b^2(2bc^2-3a^3)$

Algebra tiles could be used to factor the binomials in parts a and b

12. a) $3(4+2g-g^2)$ b) $d(3c^2-10c-2)$

c) $4mn(2n-3-4m)$ d) $y(y^3-12y+24)$

e) $10x^2y(3-2y+xy)$ f) $-4b(2b^2-5b+1)$

13. a) $4x(2x-3)$ b) $3y(y^2-4y+5)$

c) $2b(2b^2-1-3b)$ d) $6m(m^2-2-4m)$

14. a) $5q(3p^2+5pq-7q^2)$ b) $-3(4mn-5m^2-6n^2)$

15. a)




b)



c) Cannot be arranged as a rectangle

d)



16. a) Cannot be arranged as a rectangle

b)



c)



d) Cannot be arranged as a rectangle

17. 6 x-tiles

18. a) $g^2 + g - 20$

	g	-4
g	$(g)(g) = g^2$	$(g)(-4) = -4g$
5	$(5)(g) = 5g$	$(5)(-4) = -20$

b) $h^2 + 14h + 49$

	h	7
h	$(h)(h) = h^2$	$(h)(7) = 7h$
7	$(7)(h) = 7h$	$(7)(7) = 49$

c) $k^2 + 7k - 44$

	k	11
k	$(k)(k) = k^2$	$(k)(11) = 11k$
-4	$(-4)(k) = -4k$	$(-4)(11) = -44$

d) $81 - s^2$

	9	$-s$
9	$(9)(9) = 81$	$(9)(-s) = -9s$
s	$(s)(9) = 9s$	$(s)(-s) = -s^2$

e) $144 - 24t + t^2$

	12	$-t$
12	$(12)(12) = 144$	$(12)(-t) = -12t$
$-t$	$(-t)(12) = -12t$	$(-t)(-t) = t^2$

f) $42 - r - r^2$

	6	$-r$
7	$(7)(6) = 42$	$(7)(-r) = -7r$
r	$(r)(6) = 6r$	$(r)(-r) = -r^2$

g) $y^2 - 14y + 33$

	y	-11
y	$(y)(y) = y^2$	$(y)(-11) = -11y$
-3	$(-3)(y) = -3y$	$(-3)(-11) = 33$

h) $x^2 - 25$

	x	5
x	$(x)(x) = x^2$	$(x)(5) = 5x$
-5	$(-5)(x) = -5x$	$(-5)(5) = -25$


19. a) $(q+2)(q+4)$ b) $(n+5)(n-9)$
 c) $(6-s)(9-s)$ d) $(k+6)(k-15)$
 e) $(x+4)(x-5)$ f) $(3-y)(4-y)$
 20. a) i) $(m+3)(m+4)$ ii) $(m+2)(m+6)$
 iii) $(m+1)(m+12)$ iv) $(m-3)(m-4)$
 v) $(m-2)(m-6)$ vi) $(m-1)(m-12)$

b) No


21. a) $(u-3)(u-9)$ b) $(v+4)(v-5)$

c) $(w-2)(w+12)$

22. a) $2h^2 + 10h + 8$




b) $3j^2 + 16j + 5$



c) $6k^2 + 7k + 2$



d) $8m^2 + 14m + 3$



23. a) i) $2x^2 + 5x + 3$

ii)



iii) $(x+1)(2x+3)$

b) i) $3x^2 + 10x + 8$

ii)



iii) $(x+2)(3x+4)$

24. a) $6r^2 + 31r + 35$

	$3r$	5
$2r$	$(2r)(3r) = 6r^2$	$(2r)(5) = 10r$
7	$(7)(3r) = 21r$	$(7)(5) = 35$

b) $9y^2 - 80y - 9$

	y	-9
$9y$	$(9y)(y) = 9y^2$	$(9y)(-9) = -81y$
1	$(1)(y) = y$	$(1)(-9) = -9$

c) $4a^2 - 26a + 42$

	$2a$	-6
$2a$	$(2a)(2a) = 4a^2$	$(2a)(-6) = -12a$
-7	$(-7)(2a) = -14a$	$(-7)(-6) = 42$

d) $9w^2 - 9w + 2$

	$3w$	-1
$3w$	$(3w)(3w) = 9w^2$	$(3w)(-1) = -3w$
-2	$(-2)(3w) = -6w$	$(-2)(-1) = 2$

e) $16p^2 + 40p + 25$

	$4p$	5
$4p$	$(4p)(4p) = 16p^2$	$(4p)(5) = 20p$
5	$(5)(4p) = 20p$	$(5)(5) = 25$

f) $3y^2 - 2y - 1$

	$-3y$	-1
$-y$	$(-y)(-3y) = 3y^2$	$(-y)(-1) = y$
1	$(1)(-3y) = -3y$	$(1)(-1) = -1$


25. a) $(k-1)(4k-3)$
 b) $(3c+1)(2c-5)$
 c) $(b-2)(4b+3)$
 d) $(a-5)(6a-1)$
 e) $(4x-1)(7x+4)$
 f) $(3x+2)(7x-2)$
26. a) $(2m-3)(3m+7)$
 b) $(4n+1)(3n-5)$
 c) $(4p-5)(5p+4)$
27. a) $c^3 + 4c^2 + 5c + 2$
 b) $8r^3 - 22r^2 - 9r + 30$
 c) $-2j^3 - 5j^2 + 35j + 11$
 d) $6x^3 + 5x^2 - 17x - 6$
28. a) $16m^2 - 8mp + p^2$
 b) $9g^2 - 24gh + 16h^2$
 c) $y^2 - yz - 2z^2 - 2y + 4z$
 d) $-18c^2 + 39cd - 20d^2 + 21c - 28d$
29. a) $2m^4 + 7m^3 + 12m^2 + 17m + 10$
 b) $5 - 11x - 3x^2 + 11x^3 - 2x^4$
 c) $-6k^4 + 25k^3 + 10k^2 - 33k - 18$
 d) $3 + 2h - 10h^2 - 3h^3 + 2h^4$
30. a) $22a^2 + 3a + 7$
 b) $23c^2 - 10c - 53$

31. a) $n+2, n+4$
 b) $n(n+2)(n+4) = n^3 + 6n^2 + 8n$
32. a) $(9+2b)(9-2b)$
 b) $(4v+7)(4v-7)$
 c) $16(2g+h)(2g-h)$
 d) $2(3m+n)(3m-n)$
33. a) $(m-7)^2$
 b) $(n+5)^2$
 c) $(2p+3)^2$
 d) $(4-5q)^2$
 e) $(2r+7)^2$
 f) $(6-11s)^2$
34. a) $(g+3h)^2$
 b) $(4j-3k)^2$
 c) $(5t+2u)^2$
 d) $(3v-8w)^2$
35. $3x^2 + 14x + 16$

Chapter 3: Practice Test, page 201

1. A
 2. C
 3. 900; 5
 4. a) i) 20: 5, 20, 45, 80, 125, ...
 45: 5, 20, 45, 80, 125, ...
 50: 2, 8, 18, 32, 50, ...
 ii) 20: 50, 400, 1350, 3200, 6250, ...
 45: 75, 600, 2025, 4800, 9375, ...
 50: 20, 160, 540, 1280, 2500, ...

5. a) $6c^2 + 19c + 10$



- b) $72 + 86r + 24r^2$

	8	$6r$
9	$(9)(8) = 72$	$(9)(6r) = 54r$
$4r$	$(4r)(8) = 32r$	$(4r)(6r) = 24r^2$

c) $12t^2 + 13t - 35$
 $3t$

		7
4t	$(4t)(3t) = 12t^2$	$(4t)(7) = 28t$
-5	$(-5)(3t) = -15t$	$(-5)(7) = -35$

6. a) $2p^3 + 3p^2 - 16p + 7$
 b) $3e^3 + 6e^2f + 2ef^2 + 4f^3 + 5ef + 10f^2$
 c) $-7y^2 + 60yz - 16z^2$
 7. a) $(f+1)(f+16)$
 b) $(c-2)(c-11)$
 c) $(t+4)(4t-7)$
 d) $(2r+5s)^2$
 e) $(2x-5y)(3x-y)$
 f) $(h+5j)(h-5j)$
 8. $6r^3 + 11r^2 + 6r + 1$
 9. $8t^2 \pm 25t + 3$; $8t^2 \pm 14t + 3$; $8t^2 \pm 11t + 3$; $8t^2 \pm 10t + 3$

Chapter 4 Roots and Powers, page 202

4.1 Math Lab: Estimating Roots, page 206

1. Answers will vary. For example:
 a) $\sqrt{25}$, $\sqrt[3]{19}$, $\sqrt[4]{37}$, $\sqrt[5]{3}$
 b) For $\sqrt{25}$, the radicand is 25 and the index is 2.
 For $\sqrt[3]{19}$, the radicand is 19 and the index is 3.
 For $\sqrt[4]{37}$, the radicand is 37 and the index is 4.
 For $\sqrt[5]{3}$, the radicand is 3 and the index is 5.
 c) The index tells which root to take.
 2. a) 6 ; $36 = (6)(6)$
 b) 2 ; $8 = (2)(2)(2)$
 c) 10 ; $1000 = (10)(10)(10)(10)$
 d) -2 ; $(-2)(-2)(-2)(-2)(-2) = -32$
 e) $\frac{3}{5}$; $\left(\frac{3}{5}\right)\left(\frac{3}{5}\right)\left(\frac{3}{5}\right) = \frac{27}{125}$
 f) 1.5 ; $(1.5)(1.5) = 2.25$
 g) 0.5 ; $(0.5)(0.5)(0.5) = 0.125$
 h) 5 ; $(5)(5)(5)(5) = 625$
 3. a) 2.8 b) 2.1
 c) 1.8 d) 3.6
 e) 2.5 f) 2.0
 g) 4.4 h) 2.7

4. a) The calculator returns an error message; the square of a real number will always be positive.

b) Any non-zero even index

c) i) Any odd index

ii) Any even index

5. a) i) $\sqrt{4}$ ii) $\sqrt[3]{8}$
 iii) $\sqrt[4]{16}$
 b) i) $\sqrt{9}$ ii) $\sqrt[3]{27}$
 iii) $\sqrt[4]{81}$
 c) i) $\sqrt{16}$ ii) $\sqrt[3]{64}$
 iii) $\sqrt[4]{256}$
 d) i) $\sqrt{100}$ ii) $\sqrt[3]{1000}$
 iii) $\sqrt[4]{10\,000}$
 e) i) $\sqrt{0.81}$ ii) $\sqrt[3]{0.729}$
 iii) $\sqrt[4]{0.6561}$
 f) i) $\sqrt{0.04}$ ii) $\sqrt[3]{0.008}$
 iii) $\sqrt[4]{0.0016}$


6. Answers will vary. For example:

- a) $\sqrt[3]{216} = 6$ b) $\sqrt[3]{-343} = -7$
 c) $\sqrt[4]{\frac{81}{16}} = \frac{3}{2}$ d) $\sqrt{17} \approx 4.1$

4.2 Irrational Numbers, page 211


3. a) Irrational
 b) Rational
 c) Irrational
 d) Rational
 e) Irrational
 f) Rational
 4. a) 7 , $\sqrt[3]{27}$
 b) -5 , 7 , $\sqrt[3]{27}$
 c) $\frac{4}{3}$, $0.3\bar{4}$, -5 , -2.1538 , $\sqrt[3]{27}$, 7
 d) $\sqrt[4]{9}$
 5. a) $\sqrt{49} = 7$; $\sqrt[4]{16} = 2$
 b) $\sqrt{21}$ and $\sqrt[3]{36}$ cannot be written as a terminating or repeating decimals.
 6. a) Rational
 b) Irrational

7. a), b)




8. The cubes roots of the numbers in parts c and d will be irrational.


9. a)




b)



c)



d)



10. a) $\sqrt[3]{400}$, $\sqrt{50}$, $\sqrt[3]{70}$, $\sqrt[4]{100}$

b) $\sqrt{89}$, $\sqrt[3]{150}$, $\sqrt[4]{250}$, $\sqrt[3]{-150}$

11. $\sqrt[3]{98}$, $\sqrt{40}$, $\sqrt[3]{300}$, $\sqrt[3]{500}$, $\sqrt{75}$, $\sqrt{98}$

12. $\frac{-14}{5}$, $\sqrt[3]{-10}$, -2 , $\frac{123}{99}$, $\sqrt{4}$;

irrational: $\sqrt[3]{-10}$; rational: $\frac{-14}{5}$, -2 , $\frac{123}{99}$, $\sqrt{4}$

13. $\sqrt{5^2+3^2} = \sqrt{34}$, which is an irrational number.

14. a) i) True ii) True
iii) False iv) False
v) True


b) iii) 0 iv) π

15. Answers will vary. For example:

a) i) 0.75 ii) 0
iii) $\sqrt{7}$

16. Additional numbers may vary. For example:

a), b)




17. Answers may vary. For example:

a) 21 b) 125

18. a) 1.6

b)



19. 755:481 is approximately equivalent to 1.6:1,
and $\frac{1+\sqrt{5}}{2}$ is approximately 1.6.

20. a) Irrational number

b) Rational number

21. Each prime factor occurs a multiple of n times.

22. Triangles will vary. For example:

a) Side lengths: 3 units, 4 units, 5 units

b) Side lengths: 1 unit, $\sqrt{3}$ units, 2 units

c) Side lengths: 1 unit, $\sqrt{2}$ units, $\sqrt{3}$ units

d) Side lengths: $\sqrt{2}$ units, $\sqrt{3}$ units, $\sqrt{5}$ units

23. a) Yes

b) No

24. Take rational numbers to the 12th power.

4.3 Mixed and Entire Radicals, page 218

3.

Perfect square	Square root
1	1
4	2
9	3
16	4
25	5
36	6
49	7
64	8
81	9
100	10
121	11
144	12
169	13
196	14
225	15
256	16
289	17
324	18
361	19
400	20

4. a) $2\sqrt{2}$ b) $2\sqrt{3}$
 c) $4\sqrt{2}$ d) $5\sqrt{2}$
 e) $3\sqrt{2}$ f) $3\sqrt{3}$
 g) $4\sqrt{3}$ h) $5\sqrt{3}$
 5. a) $\sqrt{50}$ b) $\sqrt{72}$
 c) $\sqrt{98}$ d) $\sqrt{128}$
 e) $\sqrt{75}$ f) $\sqrt{108}$
 g) $\sqrt{147}$ h) $\sqrt{192}$

6. a)

Perfect cube	Cube root
1	1
8	2
27	3
64	4
125	5
216	6
343	7
512	8
729	9
1000	10

b)

Perfect fourth power	Fourth root
1	1
16	2
81	3
256	4
625	5

9. 25 is a perfect square, but neither 10 nor 5 is a perfect square.


10. a) $3\sqrt{10}$ b) Cannot be simplified
 c) $6\sqrt{3}$ d) $10\sqrt{6}$
 e) $3\sqrt{6}$ f) Cannot be simplified
 g) $2\sqrt{7}$ h) Cannot be simplified
 i) $4\sqrt{7}$
 11. a) $2\sqrt[3]{2}$ b) $3\sqrt[3]{3}$
 c) $4\sqrt[3]{4}$ d) $4\sqrt[3]{2}$
 e) Cannot be simplified f) $4\sqrt[3]{3}$
 g) $3\sqrt[3]{5}$ h) Cannot be simplified
 i) $5\sqrt[3]{4}$ j) $5\sqrt[3]{3}$

12. a) $\sqrt{18}$ b) $\sqrt{32}$
 c) $\sqrt{180}$ d) $\sqrt{150}$
 e) $\sqrt[3]{343}$ f) $\sqrt[3]{16}$
 g) $\sqrt[3]{81}$ h) $\sqrt[3]{192}$
 i) $\sqrt[3]{250}$ j) $\sqrt[3]{72}$
13. a) Yes
 b) No
14. $6\sqrt{7}$ ft.
15. $2\sqrt[3]{25}$ cm
16. $12\sqrt{6}$ in.
17. a) $2\sqrt[4]{3}$ b) $3\sqrt[4]{5}$
 c) $5\sqrt[4]{2}$ d) $2\sqrt[4]{11}$
18. a) $\sqrt[4]{3888}$ b) $\sqrt[4]{4802}$
 c) $\sqrt[4]{972}$ d) $\sqrt[4]{3072}$
19. a) $\sqrt{2}, \sqrt{3}, \sqrt{4}, \sqrt{5}, \sqrt{6}, \sqrt{7}, \sqrt{8}, \sqrt{9}, \sqrt{10}, \sqrt{11}, \sqrt{12}, \sqrt{13}, \sqrt{14}$
 b) i) The radicands start at 2 and increase by 1 each time.
 ii) $\sqrt{51}$
 iii) 30
20. $\sqrt[3]{1024}$
21. $4\sqrt{6}$
22. a) $8\sqrt{3}, 9\sqrt{2}, 4\sqrt{5}, 6\sqrt{2}, 2\sqrt{6}$
 b) $8\sqrt{3}, 6\sqrt{5}, 4\sqrt{7}, 2\sqrt{13}$
 c) $9\sqrt{2}, 3\sqrt{17}, 5\sqrt{6}, 7\sqrt{3}, \sqrt{103}$
23. a) 2, 20, 200;
 $\sqrt{4\,000\,000}, \sqrt{400\,000\,000}$
 b) 3, 30, 300;
 $\sqrt{27\,000\,000\,000}, \sqrt{27\,000\,000\,000\,000}$
 c) $2\sqrt{2}, 20\sqrt{2}, 200\sqrt{2}$;
 $\sqrt{8\,000\,000}, \sqrt{800\,000\,000}$
 d) $2\sqrt[3]{3}, 20\sqrt[3]{3}, 200\sqrt[3]{3}$;
 $\sqrt[3]{24\,000\,000\,000}, \sqrt[3]{24\,000\,000\,000\,000}$
24. $4\sqrt{2}$ cm, 32 cm^2 ; 4 cm, 16 cm^2
25. a) i) 14.142
 ii) 141.42
 b) i) 2.8284
 ii) 4.2426
 iii) 5.6568
 iv) 7.071


Chapter 4: Checkpoint 1, page 221

1. a) 9 b) -5
 c) 4 d) 3
2. a) 3.16 b) 2.47
 c) 1.73 d) 1.87
3. Neither
4. a) Irrational b) Irrational
 c) Irrational d) Rational
 e) Rational f) Irrational


5. a)




b)




c)




d)



6. a), b) Additional numbers may vary. For example:
 Real numbers




7. a)




b) $\sqrt{65}, \sqrt{32}, \sqrt[3]{72}, \sqrt[3]{50}, \sqrt[4]{100}, \sqrt[4]{60}$

8. Areas of squares may vary. For example:

a)



b)



9. a) $3\sqrt{5}$ b) $2\sqrt[3]{12}$
 c) Cannot be simplified d) $2\sqrt[4]{3}$
 e) $2\sqrt[3]{10}$ f) Cannot be simplified
11. a) $\sqrt{63}$ b) $\sqrt[3]{32}$
 c) $\sqrt{147}$ d) $\sqrt[4]{192}$
 e) $\sqrt[3]{270}$ f) $\sqrt{396}$

4.4 Fractional Exponents and Radicals, page 227

3. a) 4 b) 6
 c) 4 d) 2
 e) -3 f) -10
4. a) 10 b) 3
 c) 4 d) -2
5. a) $\sqrt[3]{36}$ b) $\sqrt{48}$
 c) $\sqrt[5]{-30}$
6. a) $39^{\frac{1}{2}}$ b) $90^{\frac{1}{4}}$
 c) $29^{\frac{1}{3}}$ d) $100^{\frac{1}{5}}$
7. a) 1 b) 2
 c) 4 d) 8
 e) 16 f) 32
8. a) $\sqrt[3]{4^2}$, or $(\sqrt[3]{4})^2$
 b) $\sqrt[5]{(-10)^3}$, or $(\sqrt[5]{-10})^3$
 c) $\sqrt{2.3^3}$, or $(\sqrt{2.3})^3$
9. $\sqrt[3]{350}$ cm, $350^{\frac{1}{3}}$ cm

10. a) $\sqrt[3]{48^2}$, or $(\sqrt[3]{48})^2$
 b) $\sqrt[3]{(-1.8)^5}$, or $(\sqrt[3]{-1.8})^5$
 c) $\sqrt{\left(\frac{3}{8}\right)^5}$, or $\left(\sqrt{\frac{3}{8}}\right)^5$
 d) $\sqrt[4]{0.75^3}$, or $(\sqrt[4]{0.75})^3$
 e) $\sqrt[5]{\left(-\frac{5}{9}\right)^2}$, or $\left(\sqrt[5]{-\frac{5}{9}}\right)^2$
 f) $\sqrt{1.25^3}$, or $(\sqrt{1.25})^3$
11. a) $3.8^{\frac{3}{2}}$, or $3.8^{1.5}$ b) $(-1.5)^{\frac{2}{3}}$
 c) $\left(\frac{9}{5}\right)^{\frac{5}{4}}$, or $\left(\frac{9}{5}\right)^{1.25}$ d) $\left(\frac{3}{8}\right)^{\frac{4}{3}}$
 e) $\left(\frac{5}{4}\right)^{\frac{3}{2}}$, or $\left(\frac{5}{4}\right)^{1.5}$ f) $(-2.5)^{\frac{3}{5}}$, or $(-2.5)^{0.6}$
12. a) 27 b) $\frac{9}{4}$
 c) 9 d) 0.216
 e) 16 f) $\frac{8}{125}$
13. a) $4^{\frac{1}{2}}$, $\sqrt{4}$ b) $16^{\frac{1}{2}}$, $\sqrt{16}$
 c) $100^{\frac{1}{2}}$, $\sqrt{100}$ d) $9^{\frac{1}{2}}$, $\sqrt{9}$
 e) $25^{\frac{1}{2}}$, $\sqrt{25}$
14. a) $(-1)^{\frac{1}{3}}$, $\sqrt[3]{-1}$ b) $8^{\frac{1}{3}}$, $\sqrt[3]{8}$
 c) $27^{\frac{1}{3}}$, $\sqrt[3]{27}$ d) $(-64)^{\frac{1}{3}}$, $\sqrt[3]{-64}$
 e) $64^{\frac{1}{3}}$, $\sqrt[3]{64}$
15. $\left(\frac{1}{4}\right)^{\frac{3}{2}}$, $\sqrt[3]{4}$, $4^{\frac{3}{2}}$, 4^2
16. a) i) 64 ii) 27
 iii) 16 iv) 5.9160...
 v) 1.331 vi) 0.8414...
 b) i, ii, iii, v
17. Approximately 76 m
 18. 2.744
 19. Approximately 1.3 m²

20. a) Approximately 93%
 b) Approximately 81%
 c) 5 h
21. Mars; period of Earth: approximately 363.8 Earth days;
 period of Mars: approximately 688.5 Earth days
22. Karen

4.5 Negative Exponents and Reciprocals, page 233

3. a) $\frac{1}{5^4} = 5^{-4}$
 b) $\left(-\frac{1}{2}\right)^{-3} = (-2)^3$
 c) $\frac{1}{3^{-2}} = 3^2$
 d) $\frac{1}{4^{-2}} = 4^2$
4. a) 16, $\frac{1}{16}$
 b) 16, $\frac{1}{16}$
 c) 6, $\frac{1}{6}$
 d) 64, $\frac{1}{64}$
5. $\frac{1}{1024}$
6. a) $\frac{1}{2^3}$
 b) $\frac{1}{3^5}$
 c) $\frac{1}{(-7)^2}$, or $\frac{1}{7^2}$
7. a) 2^2
 b) $\left(\frac{3}{2}\right)^3$
 c) $\left(-\frac{5}{6}\right)^4$, or $\left(\frac{5}{6}\right)^4$
8. a) $\frac{1}{9}$
 b) $\frac{1}{16}$
 c) $-\frac{1}{32}$
 d) 27
 e) $\frac{9}{4}$
 f) 125
9. a) $\frac{1}{2}$
 b) $\frac{10}{3}$
 c) $\frac{1}{3}$
 d) $-\frac{1}{4}$
 e) $\frac{100}{9}$
 f) $\frac{1}{4}$
 g) $\frac{1}{27}$
 h) 125

10. Answers may vary. For example:

a) 3^{-2}
 b) $25^{-\frac{1}{2}}$
 c) $\left(\frac{1}{2}\right)^{-2}$
 d) $\left(\frac{1}{-27}\right)^{-\frac{1}{3}}$

11. \$2651.56

12. $-\frac{3125}{1024}$

13. a) $\frac{1}{81}$
 b) $\frac{1}{64}$
 c) $\frac{1}{4}$
 d) $\frac{9}{4}$
 e) $\frac{8}{27}$
 f) $\frac{32}{243}$

14. \$1266.57

15. Approximately 0.19%

16. 5^{-2} ; $\frac{1}{25} > \frac{1}{32}$

17. a) The numbers at the left are divided by 2 each time. The exponents in the powers at the right decrease by 1 each time.

b) $2 = 2^1$; $1 = 2^0$; $\frac{1}{2} = 2^{-1}$; $\frac{1}{4} = 2^{-2}$; $\frac{1}{8} = 2^{-3}$

18. 3^8 , or 6561 times as great

19. a) The exponent is positive.

- b) The exponent is negative.

- c) The exponent is 0.

20. No; if the base is between 0 and 1, the power will be

greater than 1. For example: $\left(\frac{1}{2}\right)^{-1} = 2$

21. a) Approximately 2.0×10^{20} N

- b) Answers may vary depending on researched values.

For example: approximately 1.9×10^{20} N

Chapter 4: Checkpoint 2, page 236

1. a) 2
 b) 7
 c) 16
 d) $\frac{343}{27}$
 e) -32

2. a) i) $\sqrt[3]{35^2}$, or $(\sqrt[3]{35})^2$
 ii) $\sqrt{32^3}$, or $(\sqrt{32})^3$
 iii) $\sqrt[5]{(-32)^2}$, or $(\sqrt[5]{-32})^2$
 iv) $\sqrt{400^3}$, or $(\sqrt{400})^3$
 v) $\sqrt[3]{-125}$
 vi) $\sqrt[3]{\left(\frac{8}{125}\right)^2}$, or $\left(\sqrt[3]{\frac{8}{125}}\right)^2$

- b) iii) 4 iv) 8000
 v) -5 vi) $\frac{4}{25}$

3. a) $4^{\frac{1}{3}}$
 b) $9^{\frac{1}{2}}$, or $9^{0.5}$
 c) $18^{\frac{1}{4}}$, or $18^{0.25}$
 d) $10^{\frac{3}{2}}$, or $10^{1.5}$
 e) $(-10)^{\frac{2}{3}}$

4. Approximately 53 s

5. $\sqrt[3]{3}$, $3^{\frac{2}{3}}$, $(\sqrt[3]{3})^4$, $3^{\frac{3}{2}}$, $(\sqrt{3})^5$

6. $\sqrt[3]{421\,875}$ mm, $421\,875^{\frac{1}{3}}$ mm, 75 mm

7. a) $\frac{81}{16}$ b) 4
 c) $\frac{1}{100}$ d) 2
 e) 100 f) 625
8. \$4589.06

4.6 Applying the Exponent Laws, page 241

3. a) x^7 b) $\frac{1}{a^3}$
 c) b^2 d) $\frac{1}{m}$
4. a) 0.5^5 b) 0.5^{-1}
 c) 0.5^{-1} d) 0.5^5
5. a) x^2 b) $\frac{1}{x^3}$
 c) n d) $\frac{1}{a^4}$

6. a) n^6 b) $\frac{1}{z^6}$
 c) n^{12} d) $\frac{1}{c^4}$
7. a) $\left(\frac{3}{5}\right)^{12}$ b) $\left(\frac{3}{5}\right)^{-12}$
 c) $\left(\frac{3}{5}\right)^{12}$ d) $\left(-\frac{3}{5}\right)^{12}$

8. a) $\frac{a^2}{b^2}$ b) $\frac{n^6}{m^3}$
 c) $\frac{d^8}{c^8}$ d) $\frac{4b^2}{25c^2}$
 e) a^2b^2 f) n^6m^3
 g) $\frac{1}{c^{12}d^8}$ h) $\frac{x^3}{y^3}$

9. a) x ; product of powers law
 b) a^{-5} ; product of powers law
 c) b^3 ; product of powers law
 d) 1; product of powers law
 e) $\frac{1}{x^7}$; quotient of powers law
 f) s^{10} ; quotient of powers law
 g) $\frac{1}{b^5}$; quotient of powers law
 h) 1; quotient of powers law

10. a) 2.25 b) $\frac{9}{16}$
 c) 0.36 d) 1
 e) $\frac{5}{3}$ f) $-\frac{3}{8}$
 g) $\frac{1000}{343}$ h) $\frac{3}{10}$

11. a) x^3y^6 b) $\frac{a^4}{4b^4}$
 c) $\frac{1}{64m^6n^9}$ d) $\frac{16m^8n^{12}}{81}$

12. 10.6 cm

13. 251 ft.²

14. a) $\frac{a^5}{b}$ b) $\frac{d^4}{c^2}$

15. a) -32 b) $-\frac{1}{8}$
 c) $-\frac{1}{32}$ d) $\frac{1}{1024}$

16. a) m^2 b) $\frac{1}{x^{\frac{5}{4}}}$
- c) $-\frac{3b^{\frac{1}{2}}}{a^6}$ d) $-\frac{4c^2b^{\frac{1}{6}}}{a^3}$
17. a) $\frac{x^{\frac{5}{2}}}{y^4}$ b) $\frac{b}{25a^4}$
19. a) $\frac{m^8}{n^2}$ b) $\frac{r^{\frac{1}{2}}}{s^4}$
20. a) i) Dimensions, in millimetres: $\frac{1000}{2^4}$ by $\frac{1000}{2^4}$;
297 mm by 420 mm
ii) Dimensions, in millimetres: $\frac{1000}{2^4}$ by $\frac{1000}{2^4}$;
210 mm by 297 mm
iii) Dimensions, in millimetres: $\frac{1000}{2^4}$ by $\frac{1000}{2^4}$;
149 mm by 210 mm
- b) i) Dimensions, in millimetres: $\frac{1000}{2^4}$ by $\frac{1000}{2^4}$
ii) Dimensions, in millimetres: $\frac{1000}{2^4}$ by $\frac{1000}{2^4}$
iii) Dimensions, in millimetres: $\frac{1000}{2^4}$ by $\frac{1000}{2^4}$
- c) A piece of A4 paper has the same dimensions as a folded piece of A3 paper; a piece of A5 paper has the same dimensions as a folded piece of A4 paper.
21. a) $\frac{a^{16}c^3}{b^7}$ b) $\frac{c^{14}}{64a^2b^{10}}$
22. a) $\frac{1}{a^{\frac{10}{9}}}$ b) $\frac{1}{a^{\frac{7}{2}}}$
23. For example:
- a) $x^1 \cdot x^{\frac{1}{2}}, x^{\frac{3}{4}} \cdot x^{\frac{3}{4}}, x^2 \cdot x^{-\frac{1}{2}}$
- b) $x^2 \div x^{\frac{1}{2}}, x^{\frac{5}{2}} \div x^1, x^{-1} \div x^{-\frac{5}{2}}$
- c) $\left(x^{\frac{1}{2}}\right)^3, \left(x^6\right)^{\frac{1}{4}}, \left(x^{-\frac{1}{3}}\right)^{-\frac{9}{2}}$
24. $\frac{1}{2}\left(\frac{3}{2}\right)^{\frac{1}{2}}$ cm, or approximately 0.6 cm


Chapter 4: Review, page 246

1. a) 10 b) 0.9
c) 2 d) $\frac{3}{5}$
2. The index tells which root to take.
3. a) 3.3 b) -2.3
c) 2.0
4. a) 25 b) 216
c) 2401
5. Neither
6. a) Rational b) Rational
c) Rational d) Irrational
e) Rational f) Rational
g) Rational h) Irrational
i) Irrational
7. Approximately 4.8 cm
8. a) Rational b) Irrational
9. $\sqrt[3]{-30}, \sqrt[4]{10}, \sqrt[4]{18}, \sqrt[3]{30}, \sqrt{20}, \sqrt{30}$
-
10. 1 s
11. a) $5\sqrt{6}$ b) $3\sqrt[3]{5}$
c) $4\sqrt{7}$ d) $3\sqrt[4]{2}$
12. a) $\sqrt{180}$ b) $\sqrt{126}$
c) $\sqrt[3]{192}$ d) $\sqrt[4]{32}$
13. Approximately 1.0 cm
15. $6\sqrt{2}, 3\sqrt{6}, 5\sqrt{2}, 4\sqrt{3}, 2\sqrt{7}$
17. a) $\sqrt[4]{12}$ b) $\sqrt[3]{(-50)^5}$, or $(\sqrt[3]{-50})^5$
c) $\sqrt{1.2}$ d) $\sqrt[3]{\frac{3}{8}}$
18. a) $1.4^{\frac{1}{2}}$ b) $13^{\frac{2}{3}}$
c) $2.5^{\frac{4}{5}}$ d) $\left(\frac{2}{5}\right)^{\frac{3}{4}}$
19. a) 2 b) 1.2
c) -32 d) $\frac{27}{64}$
20. Approximately 35%
21. $(\sqrt{5})^3, 5^{\frac{3}{4}}, 5^{\frac{2}{3}}, \sqrt[3]{5}, \sqrt[4]{5}$

22. a) Approximately 7122 Calories/day
b) Approximately 4 Calories/day
23. a) The numbers at the left are divided by 3 each time; the exponents in the powers at the right decrease by 1 each time.
b) $3 = 3^1$; $1 = 3^0$; $\frac{1}{3} = 3^{-1}$; $\frac{1}{9} = 3^{-2}$; $\frac{1}{27} = 3^{-3}$
24. a) $\frac{1}{4}$
b) $\frac{27}{8}$
c) $\frac{125}{8}$
25. \$908.51
26. 18.0 cm
27. 262 Hz
28. a) $9m^8n^2$
b) $\frac{1}{x^4y^6}$
c) $\frac{1}{4ab^3}$
d) $\frac{1}{r^{\frac{10}{3}}s^{\frac{2}{3}}}$
29. a) a^2b^5
b) $\frac{x^2}{y}$
c) $\frac{1}{a^5}$
d) $x^{\frac{3}{2}}y^3$
30. a) $\frac{9}{4}$
b) 30.25
c) $\frac{144}{25}$
d) 0.4
31. Approximately 6.4 cm
32. a) $s^3t^{\frac{10}{3}}$
b) $\frac{d^9}{64c}$

Chapter 4: Practice Test, page 249

1. B
2. A
3. a) $5\sqrt{3}$; $5\sqrt{3} = \sqrt{75}$
b)



4. a) $\frac{4}{3}$
b) $\frac{1}{16}$
c) 0.729
d) $\frac{1}{4}$

5. $2\sqrt{11}$
6. $\frac{y^5}{x^2}$
7. a) $\frac{1}{p^2q}$
b) $\frac{1}{cd^{\frac{1}{3}}}$
8. Approximately 29 L

Cumulative Review Chapters 1–4, page 252

1. 117 m^2
2. 236 in.^3
3. a) 5.2 cm
b) 1 in.
4. 28 ft.
5. 64.2°
6. a) $9\frac{7}{10} \text{ in.}$
b) 4 in.^2
7. a) 9; 585
b) 14; 924
c) 3; 3150
d) 2; 4620
8. 8214 in.^2
9. a) 1, 4, 9, 16, 25, 36, 49, 64, 81, 100
b) 1, 8, 27, 64, 125, 216, 343, 512, 729, 1000
c) 1, 64, 729
10. a) $3a(5a - 9)$
b) $2p(2 + 6p^2 - 3p)$
c) $-2d(4d^3 + 7)$
d) $7(3w - 4 + 2w^2)$
e) $2x^2y^2(9x^2 - 2xy + 5y^2)$
f) $11np^2(3n^3p + n - 11p^2)$
11. The trinomials that can be represented as a rectangle of algebra tiles can be factored.
a) Can be represented
b) Cannot be represented
c) Cannot be represented
d) Can be represented
12. a) $d^2 + 2d - 15$

	d	-3
d	$(d)(d) = d^2$	$(d)(-3) = -3d$
5	$(5)(d) = 5d$	$(5)(-3) = -15$

- b) $45 - 14s + s^2$

	9	$-s$
5	$(5)(9) = 45$	$(5)(-s) = -5s$
$-s$	$(-s)(9) = -9s$	$(-s)(-s) = s^2$

c) $-49 + 16g^2$

	7	$4g$
-7	$(-7)(7) = -49$	$(-7)(4g) = -28g$
$4g$	$(4g)(7) = 28g$	$(4g)(4g) = 16g^2$

d) $6k^2 + 13k - 63$

	$2k$	9
$3k$	$(3k)(2k) = 6k^2$	$(3k)(9) = 27k$
-7	$(-7)(2k) = -14k$	$(-7)(9) = -63$

13. Answers may vary. For example, one of these:

- a) 15, -15, 9, -9
- b) 6, 4, 0, -6, -14, -24, -36, ...
- c) 17, -17, 7, -7, 3, -3
- d) 4, 3, 0, -5, -12, -21, -32, ...

14. a) $(n + 11)(n - 2)$

b) $(4 - m)(15 - m)$

c) $(2r + 5)(3r + 4)$

d) $(2n + 1)(5n - 2)$

15. a) $3(c - 10)(c + 2)$

b) $-5(h + 7)(h - 3)$

c) $3(8c + 3)(c - 4)$

d) $5(4 - 3a)(5 - 4a)$

e) $4(t - 6)^2$

f) $2(4 + w)(8 - w)$

g) $3(6r - 7s)(6r + 7s)$

h) $-2(5x - 3y)(7x + 2y)$

16. a) $2x^3 + 3x^2 - 19x + 15$

b) $2a^2 - ab - 6a - 10b^2 - 12b$

c) $12 - t - t^2 + 9s - 3st$

d) $2n^4 + 3n^3 - 8n^2 - 7n + 4$

17. a) $5c^2 + 23c - 42$

b) $-2t^2 - 33t + 30$

c) $-4w^2 + 53w + 46$

d) $3d^2 + 12d - 25$

18. a) $(5n + 4)^2$


b) $(6v - w)(4v + 3w)$

c) $(9c - 13d)(9c + 13d)$

d) $(3a - 5b)^2$

19. 3.42

20. $\sqrt[3]{-90}$, $\sqrt[4]{150}$, $\sqrt[4]{250}$, $\sqrt[3]{90}$, $\sqrt{30}$



21. a) i) $4\sqrt{6}$

ii) $3\sqrt[3]{4}$

iii) $2\sqrt[4]{9}$

iv) $5\sqrt{17}$

v) $6\sqrt[3]{3}$

vi) $2\sqrt[4]{22}$

b) i) $\sqrt{75}$

ii) $\sqrt[3]{40}$

iii) $\sqrt[4]{29 \cdot 282}$

iv) $\sqrt{63}$

v) $\sqrt[3]{2916}$

vi) $\sqrt[3]{96}$

22. a) i) $\sqrt[4]{50^3}$, or $(\sqrt[4]{50})^3$

ii) $\sqrt[3]{(-2.5)^2}$, or $(\sqrt[3]{-2.5})^2$

iii) $\sqrt[5]{\left(\frac{3}{4}\right)^8}$, or $\left(\sqrt[5]{\frac{3}{4}}\right)^8$

b) i) $8.9^{\frac{2}{3}}$

ii) $\left(\frac{7}{4}\right)^{\frac{3}{4}}$

iii) $(-4.8)^{\frac{6}{5}}$

23. a) 27

b) $\frac{216}{343}$

c) -0.002 43

d) $\frac{81}{16}$

e) $\frac{1}{8}$

f) $\frac{512}{125}$

g) 27

h) $\frac{25}{4}$, or 6.25

i) $\frac{1331}{343}$

24. \$24 895.92

25. a) $\frac{4}{25}$

b) 0.25

c) $\frac{5}{3}$

d) $-\frac{1}{2}$

26. a) a^3b^2

b) $\frac{16x^{24}}{y^8}$

c) $\frac{-3b^{\frac{5}{2}}}{a^{\frac{3}{2}}}$

d) $\frac{-5z}{x^2y^3}$


5.1 Representing Relations, page 262

3. a) i) The relation shows the association “has a value, in dollars, of” from a set of coins to a set of numbers.

ii) As a set of ordered pairs:

{(penny, 0.01), (nickel, 0.05), (dime, 0.10), (quarter, 0.25), (loonie, 1.00), (toonie, 2.00)}

As an arrow diagram:




- b) i) The relation shows the association “is played with a” from a set of sports to a set of equipment.

ii) As a set of ordered pairs:

{(badminton, racquet), (badminton, shuttlecock), (hockey, puck), (hockey, stick), (tennis, ball), (tennis, racquet), (soccer, ball)}


As an arrow diagram:



4. a) As a table:

Word	Number of Letters
blue	4
green	5
orange	6
red	3
yellow	6

- b) As an arrow diagram:




5. a) The relation shows the association “creates art using the medium of” from a set of francophone artists from Manitoba to a set of artistic mediums.

b) i) As a set of ordered pairs:

{(Gaëtan Sylvester, sculpture), (Hubert Théroux, painting), (Huguette Gauthier, stained glass), (James Culleton, painting), (Nathalie Dupont, photography), (Simone Hébert Allard, photography)}

ii) As an arrow diagram:




6. a) The relation shows the association “has a typical mass, in kilograms, of” from a set of salmon species to a set of masses.

b) As a set of ordered pairs:

{(Chinook, 13), (Chum, 9), (Coho, 5), (Pink, 3), (Sockeye, 6)}

c) As an arrow diagram:



7. a) The arrow diagram shows a relation with the association “is the number of letters in” from a set of numbers to a set of words beginning with the letter Z.

b) As a set of ordered pairs:


{(3, Zen), (4, zany), (4, zero), (5, zebra), (6, zombie), (7, Zamboni), (8, zeppelin)}

As a table:

Number	Word beginning with Z
3	Zen
4	zany
4	zero
5	zebra
6	zombie
7	Zamboni
8	zeppelin

- c) Chosen words and representations may vary. For example:

As an arrow diagram:



As a set of ordered pairs:

{(4, X-ray), (5, xenon), (5, Xerox), (5, xylem), (9, xylophone), (10, xenophilia)}

As a table:

Number	Word beginning with X
4	X-ray
5	xenon
5	Xerox
5	xylem
9	xylophone
10	xenophilia


8. a) The diagram shows a relation with the association “translates to” from the set of French words to the set of English words.

b) Answers may vary. For example: Two ordered pairs that satisfy the relation are: (oui, yes) and (et, and)

9. a) {(0, 6), (1, 2), (2, 5), (3, 5), (4, 4), (5, 5), (6, 6), (7, 3), (8, 7), (9, 6)} Some digital clocks may show the number 9 with 5 line segments.

b) Representations may vary. For example:


As an arrow diagram:



As a table of values:

Digit	Number of lit segments
0	6
1	2
2	5
3	5
4	4
5	5
6	6
7	3
8	7
9	6

10. a) As an arrow diagram:




As a set of ordered pairs:

{(Hayley Wickenheiser, 1978), (Jennifer Botterill, 1979), (Jonathan Cheechoo, 1980), (Jordin Tootoo, 1983), (Roberto Luongo, 1979)}

As a table:

Hockey Player	Birth Year
Hayley Wickenheiser	1978
Jennifer Botterill	1979
Jonathan Cheechoo	1980
Jordin Tootoo	1983
Roberto Luongo	1979

b) As an arrow diagram:



As a set of ordered pairs:

$\{(1978, \text{Hayley Wickenheiser}),$
 $(1979, \text{Jennifer Botterill}),$
 $(1979, \text{Roberto Luongo}),$
 $(1980, \text{Jonathan Cheechoo}),$
 $(1983, \text{Jordin Tootoo})\}$

As a table:

Birth Year	Hockey Player
1978	Hayley Wickenheiser
1979	Jennifer Botterill
1979	Roberto Luongo
1980	Jonathan Cheechoo
1983	Jordin Tootoo

11. Answers may vary. For example:

- a) Ordered pairs should be in the form: (older person, younger person)
 b) Other associations include: “is taller than”
 “is involved in more school groups than”
 “usually wakes up earlier than”

12. a) i) $\{(1, 1), (1, 3), (1, 5), (2, 2), (2, 4), (2, 6), (3, 1), (3, 3), (3, 5), (4, 2), (4, 4), (4, 6), (5, 1), (5, 3), (5, 5), (6, 2), (6, 4), (6, 6)\}$
 ii) $\{(1, 3), (1, 4), (1, 6), (2, 4), (2, 5), (3, 1), (3, 5), (3, 6), (4, 1), (4, 2), (4, 6), (5, 2), (5, 3), (6, 1), (6, 3), (6, 4)\}$

b) No

13. a) 6 children b) 4 parents

c) 2 grandparents

14. a) 2 females

b) 3 males

5.2 Properties of Functions, page 270

4. a) Function

b) Not a function

c) Function

5. a) Function; domain: $\{1, 2, 3, 4\}$; range: $\{3, 6, 9, 12\}$

b) Not a function; domain: $\{-1, 0, 1\}$; range: $\{-1, 0, 1\}$

c) Function; domain: $\{2, 4, 6, 8\}$; range: $\{3, 5, 7, 9\}$

d) Not a function; domain: $\{0, 1, 2\}$; range: $\{1, 2, 3\}$

6. a) $C(n) = 20n + 8$

b) $P(n) = n - 3$

c) $t(d) = 5d$

d) $f(x) = -x$

7. a) $d = 3t - 5$

b) $y = -6x + 4$

c) $C = 5n$

d) $P = 2n - 7$

8. a) Function; domain: $\{1, 2, 3, 4\}$; range: $\{1, 8, 27, 64\}$

b) Not a function; domain: $\{3\}$; range: $\{4, 5, 6, 7\}$

9. a) i) Function

ii) Dependent variable: C ; independent variable: n

iii) Domain: $\{1, 2, 3, 4, 5, 6, \dots\}$;

range: $\{2.39, 4.00, 6.39, 8.00, 10.39, 12.00, \dots\}$

b) i) Function

ii) Dependent variable: T ; independent variable: A

iii) Domain: $\{610, 1220, 1830, 2440, 3050, 3660, \dots\}$;

range: $\{15.0, 11.1, 7.1, 3.1, -0.8, -4.8, \dots\}$

10. a) Not a function

b) Function

c) Part a: domain: $\{3, 4, 5, 6\}$; range: $\{\text{equilateral triangle, hexagon, isosceles triangle, parallelogram, pentagon, rectangle, rhombus, right triangle, scalene triangle, square, trapezoid}\}$

Part b: domain: $\{\text{equilateral triangle, hexagon, isosceles triangle, parallelogram, pentagon, rectangle, rhombus, right triangle, scalene triangle, square, trapezoid}\}$; range: $\{3, 4, 5, 6\}$

11. Answers may vary. For example:

a) Functions:

Name	From
Marie	Edmonton
Gabriel	Falher
Élise	Bonnyville
Christophe	Calgary
Jean	Edmonton
Mélanie	Edmonton
Nicole	Red Deer
Marc	Légal

Name	Age
Marie	13
Gabriel	16
Élise	14
Christophe	13
Jean	15
Mélanie	15
Nicole	17
Marc	13

b) Not functions:

Age	Name
13	Marie
16	Gabriel
14	Élise
13	Christophe
15	Jean
15	Mélanie
17	Nicole
13	Marc

From	Age
Edmonton	13
Falher	16
Bonnyville	14
Calgary	13
Edmonton	15
Edmonton	15
Red Deer	17
Légal	13

12. The statement in part a is true.

13. a)

Letter	Number
A	1
D	2
F	4
G	2
M	3
Q	10
T	1
X	8
Z	10

Number	Letter
1	A
1	T
2	D
2	G
3	M
4	F
8	X
10	Q
10	Z

b) The first table represents a function.

14. a) $f(1) = 6$

b) $f(-3) = 26$

c) $f(0) = 11$

d) $f(1.2) = 5$

15. a) i) $n = 9$

ii) $n = \frac{1}{2}$, or 0.5

b) i) $x = -8$

ii) $x = \frac{17}{5}$, or 3.4

16. a) $C = 2.54i$

b) $C(12) = 30.48$; a length of 12 in. is equal to a length of 30.48 cm.

c) $i = 39.3700\dots$; a length of 100 cm is approximately equal to a length of 39 in.

17. a) $D(t) = -80t + 300$

b) 300 km

18. a) i) $f(15) = 112.785$; a female whose humerus is 15 cm long will be approximately 113 cm tall.

ii) $m(20) = 128.521$; a male whose humerus is 20 cm long will be approximately 129 cm tall.

b) i) $l = 25.6082\dots$; a female who is 142 cm tall will have a humerus length of approximately 26 cm.

ii) $l = 42.6257\dots$; a male who is 194 cm tall will have a humerus length of approximately 43 cm.

19. a) i) $C(50) = 10$

ii) $C(-13) = -25$

b) i) $f = 68$

ii) $f = -31$

c) i) $C(32) = 0$

ii) $C(212) = 100$

iii) $C(356) = 180$

20. Variables may differ. Let c represent a temperature in degrees Celsius. Let F represent the same temperature in degrees Fahrenheit. $F(c) = \frac{9}{5}c + 32$

21. $P(l) = 2l + \frac{18}{l}$

22. $l(w) = 6 - w$; domain: $0 < w < 6$; range: $0 < l < 6$


23. $t(s) = 11 - 2s$; domain: $1.5 < s < 3$; range: $5 < t < 8$

Chapter 5: Checkpoint 1, page 275

1. a) In words:

This relation shows the association “is used to make” from a set of materials to a set of objects.

As an arrow diagram:



As a table:

Material	Object
bark	basket
skin	drum
skin	kayak
stone	carving
stone	inukshuk


b) In words:

This relation shows the association “has this many factors” from the natural numbers from 1 to 4 to a set of natural numbers.

As a set of ordered pairs:

$\{(1, 1), (2, 2), (3, 2), (4, 3)\}$

As an arrow diagram:
has this many factors



c) In words:

This relation shows the association “is usually coloured” from a set of objects to a set of colours.

As a set of ordered pairs:

{(grass, green), (sea, blue), (sky, blue), (snow, white)}


As a table:

Object	Colour
grass	green
sea	blue
sky	blue
snow	white

d) As a set of ordered pairs:

{(2, 1), (3, 1), (3, 2), (4, 1), (4, 2), (4, 3)}

As an arrow diagram:
is greater than



As a table:

Number	Number
2	1
3	1
3	2
4	1
4	2
4	3

2. a) The relations in parts b and c are functions.

b) Part b: domain: {1, 2, 3, 4}; range: {1, 2, 3}


Part c: domain: {grass, sea, sky, snow};
range: {blue, green, white}

3. Answers may vary. For example:

a) i) {(1, 1), (1, 3), (1, 5), (1, 7)}

ii) {(1, 1), (3, 3), (5, 5), (7, 7)}


b) i) As an arrow diagram:
is less than or equal to



As a table of values:

Number	Number
1	1
1	3
1	5
1	7

ii) As an arrow diagram:
plus 0 is



As a table of values:

Number	Number
1	1
3	3
5	5
7	7

4. a) Dependent variable: T ; independent variable: d

b) $T = 10d + 20$

c) $T(5) = 70$; At a depth of 5 km below Earth's surface, the temperature is 70°C .

d) $d = 3$; A temperature of 50°C occurs at a depth of 3 km below Earth's surface.

5.3 Interpreting and Sketching Graphs, page 281

3. a) Bear F; approximately 650 kg

b) Bear A; approximately 0.7 m

c) Bears D and E; 400 kg

d) Bears D and H; approximately 2.25 m

4. a) 8 m; 06:00 and 18:00

b) 2 m; 00:00 (midnight), 12:00 (noon), and 24:00 (midnight)

c) Approximately 6.5 m

d) At approximately 02:20, 09:40, 14:20, and 21:40


5. Graph B

8. a) True b) False
 c) True d) False
 e) False

9. b) 25 L; no


10.

Temperature of an Oven



12.

Number of Cartons in the School Vending Machine




13. From 3 min to 4 min, the volume should be below 40 because Jonah turns the volume down.
 At 9 min, the graph should be a vertical line from 80 to 0 because the mute button immediately silences the television.

14. Answers may vary. For example:

a)


Distance from Home



Situation: A person walks from home to a park 1.5 km away in 10 min. He sits on a park bench and reads for 10 min. Then he walks home.

b)

Speed while Sprinting




Situation: A person sprints down a street starting from a standstill. It takes the person 5 s to reach a speed of 7.5 m/s. After 5 s of running at 7.5 m/s, the person slows down and stops in 5 s.

15. Answers may vary. For example:

a)


Emptying a Watering Can



Situation: A watering can contains 4 L of water. The water is poured at a steady rate so the watering can is empty after 30 s.

b)

Height of a Helium Balloon




Situation: A person lets go of a helium balloon. The balloon starts at a height of 2.5 m above the ground. After 10 s, it is at a height of 15 m above the ground.


16. Answers may vary. For example:

a) i)

Distance above the Ground while Bungee Jumping




ii) **Speed while Bungee Jumping**



17. a)


Height of a Jump



Situation: The height of a grasshopper during one hop. It takes 0.2 s for a grasshopper to jump 20 cm high, and another 0.2 s for it to return to the ground.

b)


Cost of Parking in a Parking Garage



Situation: The cost of parking in a parking garage. It costs \$1 to park for up to 30 min, \$2 to park from 30 min to 60 min, and \$3 to park from 60 min to 90 min.

c)


Height of a Point on the Rim of a Tire



Situation: The height of a point on the rim of a tire on a truck over time. The point starts at the lowest point on the rim, 25 cm above the ground. As the wheel goes around, the point moves up to a maximum height of 50 cm, then down, then up again.


18.

Depth of Water in Two Pools




5.4 Math Lab: Graphing Data, page 286

1. a) i) The points are joined because air temperature and speed can have any numerical value between those indicated by the points on the graph.




ii) Yes

- b) i) The points are not joined because the data are only valid for whole numbers of years.




ii) Yes

2. a) The points are not joined because only whole numbers are permissible for the number of juice cans purchased.



The relation is a function because there is only one cost for each number of cans.

- b) The points are joined because all values of altitude and temperature are permissible between those plotted.




The relation is a function because there is only one value of temperature for each altitude.

5.5 Graphs of Relations and Functions, page 294

4. a) Domain: $\{-2, -1, 0, 1, 2\}$; range: $\{-4, -2, 0, 2, 4\}$
 b) Domain: $\{-3, -1, 0, 2, 3\}$; range: $\{-2, 0, 1, 2, 3\}$
 c) Domain: $\{-3, -2, -1, 0, 1, 2, 3\}$; range: $\{2\}$
5. A vertical line drawn on each graph intersects the graph at 0 points or 1 point.
6. a) Yes; each point on the line has a different x -coordinate.
 b) No; each point on the line has the same x -coordinate, 1.
7. a) iv b) i
 c) ii d) iii
8. a) Function; domain: all real numbers; range: $1 \leq y \leq 3$
 b) Not a function; domain: $-3 \leq x \leq 1$; range: $y \geq -1$
 c) Not a function; domain: $\{1, 2, 3, 4, 5\}$; range: $\{2, 3, 4, 5\}$
 d) Function; domain: $x \geq -2$; range: $2 \leq y \leq 4$
 e) Not a function; domain: $x \leq 2$; range: $1 \leq y \leq 5$
9. a) Domain: all real numbers; range: $y \geq 1$
 b) Domain: $-3 \leq x \leq 3$; range: $0 \leq y \leq 3$
 c) Domain: $-3 \leq x \leq 3$; range: $-3 \leq y \leq 0$
 d) Domain: $-1 \leq x \leq 2$; range: $0 \leq y \leq 3$
10. a) The points on the graph should not be connected.
 b) The points on the graph should be connected.
 c) The points on the graph should be connected.
 d) The points on the graph should be connected.
11. a) i) The distance of a school bus from the school from 8:00 to 9:00.
 ii) The number of students on a school bus from 8:00 to 9:00.
 b) i) Independent variable: time; dependent variable: distance from the school
 ii) Independent variable: time; dependent variable: number of students
 c) Graph A: points are connected because all values of time and distance are permissible between the indicated plotted points.
 Graph B: points are not connected because it is impossible to have only part of a student on a bus.
12. a) The points on the graph are connected because the car's speed and skid length can be any positive number of kilometres per hour and metres, respectively, between the plotted points.
 b) Exact numbers for the range may vary. For example: domain: $40 \leq s \leq 120$; range: $16 \leq d \leq 144$
 Restrictions: the domain and range cannot contain negative numbers because it is impossible to have a negative skid distance or a negative speed.
 The domain is also restricted because the relationship shown on the graph may not be true for speeds less than 40 km/h and greater than 120 km/h.
13. a) Independent variable: t ; dependent variable: n
 b) The points are not connected because it is impossible to have part of a car in a parking lot.
 c) Exact numbers for the range may vary. For example: domain: $\{8:00, 10:00, 12:00, 14:00, 16:00\}$; range: $\{4, 25, 31, 64, 65\}$
 Restrictions: the domain can be any time between 00:00 and 24:00, all the possible times in one day.
 The range can be any whole number up to the number of parking spaces in the lot.
14. a) Independent variable: number of days after January 1; dependent variable: number of hours the sun is above the horizon, h

b) Answers may vary. For example:

Number of Hours the Sun Is above the Horizon in Paulatuq



I connected the points because the relationship shown on the graph is true for days represented by points between the ones plotted.

The data are discrete, but the scale is so small that if all the points were plotted, they would make a line segment.

c) From the table: the relation is a function because each number in the first column is different.


From the graph: the relation is a function because a vertical line drawn on the graph would intersect the graph in only 1 point.

15. a)

Volume of Paint, p (L)	0	2	4	6	8
Cost, c (\$)	0	24	48	72	96
Area Covered, A (m^2)	0	17	34	51	68


b)

Area Covered by Paint



c)

Area that Can Be Covered for a Given Cost



d) Part b: domain: $0 \leq p \leq 8$; range: $0 \leq A \leq 68$

Part c: domain: $0 \leq c \leq 96$; range: $0 \leq A \leq 68$


16. a) -1

b) 3

17. a) 5


b) 3

18. Answers may vary. For example:




Domain: $-2 \leq x \leq 4$; range: $-2 \leq y \leq 4$

19. a)




b)



20. a)

Planetary Years as a Function of Distance from the Sun




The points are not joined because each point represents a planet and their number is limited.

b) Domain: {1, 5, 10, 19}; range: {1, 12, 29, 84}

21. a)

Cost of Sending a Letter in 2009



b) Domain: all real numbers greater than 0 and less than or equal to 500; range: {0.54, 0.98, 1.18, 1.96, 2.75}

22. Yes

23. The statement is false.

24. a)

Payment Scheme 1	
Day	Total money received (\$)
1	0.01
2	0.03
3	0.07
4	0.15


Payment Scheme 2	
Day	Total money received (\$)
1	10
2	20
3	30
4	40

5	0.31
6	0.63
7	1.27
8	2.55
9	5.11
10	10.23
11	20.47
12	40.95
13	81.91
14	163.83
15	327.67

5	50
6	60
7	70
8	80
9	90
10	100
11	110
12	120
13	130
14	140
15	150

b)


Total Money Received Under Two Payment Schemes




c) I would choose Payment Scheme 1 because after 13 days, the money received is greater and increases at a faster rate.

Chapter 5: Checkpoint 2, page 299

1. Answers may vary. For example:
Paula's Distance from Home



2. a)



b) No

c) Domain: {14, 15, 17, 18};
range: {45, 50, 56, 64, 90}

3. a) Not a function; domain: $0 \leq x \leq 2$; range: $1 \leq y \leq 5$

b) Function; domain: $x \geq -3$; range: $y \geq 0$

c) Function; domain: $-2 \leq x \leq 2$; range: $-8 \leq y \leq 8$

5.6 Properties of Linear Relations, page 308

3. a) Linear relation

b) Not a linear relation

c) Linear relation

d) Not a linear relation

4. a) Linear relation

b) Not a linear relation

c) Not a linear relation

5. a) Linear relation

b) Linear relation


c) Not a linear relation

d) Not a linear relation

6. a) Tables of values may vary. For example:


i)

x	y
-2	4
-1	6
0	8
1	10
2	12




ii)

x	y
-2	11
-1	11.5
0	12
1	12.5
2	13




iii)

x	y
-2	12
-1	9
0	8
1	9
2	12




iv)

x	y
-2	-4
-1	-2
0	0
1	2
2	4




v)



vi)

x	y
-2	8
0	6
2	4
4	2
6	0



b) The relations in part a, i, ii, iv, v, and vi are straight lines, so they are linear relations.

7. a) i) Independent variable: s ; dependent variable: d

ii) Not linear


b) i) Independent variable: t ; dependent variable: a

ii) Linear

iii) -200 m/min

8. a)

Distance to the Horizon for a Given Height in a Hot-Air Balloon



b) The relation is not linear because the points on the graph do not lie on a straight line.

9. Answers may vary. For example:

I could examine the change in the first and second coordinates. If both changes are constant, the relation is linear.

I could also graph the ordered pairs. If the points lie on a straight line, the relation is linear.

10. Yes

11. The first set of ordered pairs does not represent a linear relation. The second set of ordered pairs represents a linear relation.

12. a) Answers may vary. For example: The equation relates the dependent variable, C , to the rate of change, 15, times the independent variable, n , plus a constant, 550.

b) 15; cost per guest

13. Answers may vary. For example:

Create a table of values for the relation. Then, either check the differences in the numbers in each column or plot the points. If the differences are constant or the points lie along a line, the relation is linear. Otherwise, it is not linear.

14. a) Independent variable: t ; dependent variable: C

b) \$0.08/min; every minute, the cost of the phone call increases by \$0.08.

15. -\$0.80/booth; at every toll booth, Kashala pays \$0.80.

16. a) Equation 3 and Set B

b) Equation 1 and Set C

c) Equation 2 and Set A

17. a) i) Linear ii) Not linear

iii) Linear iv) Linear

v) Not linear

b) i) Independent variable: time since the hang glider started her descent; dependent variable: hang glider's altitude; rate of change: -50 m/min; every minute, the hang glider's altitude decreases by 50 m.

iii) Independent variable: distance travelled;

dependent variable: taxi fee; rate of change:

\$2/km; every kilometre, the fee increases by \$2.

iv) Independent variable: number of yearbooks to be

printed; dependent variable: fee; rate of change:

\$5/yearbook; for every yearbook to be printed, the

fee increases by \$5.

18. a) Linear

b) Not linear

c) Not linear

d) Linear

e) Not linear

19. a) The equation $V = 24\,000 - 2000n$ is linear.

The equation $V = 24\,000(0.2^n)$ is not linear.

b) -\$2000/year; every year, the value of the truck depreciates by \$2000

20. Yes; the relation is linear.

21. No; the relation is not linear.

22. a) True

b) True

c) False

d) True

e) False

5.7 Interpreting Graphs of Linear Functions, page 319

4. a) i) Vertical intercept: 0; horizontal intercept: 0; (0, 0); (0, 0)

ii) 40 km/h

iii) Domain: $0 \leq t \leq 3$; range: $0 \leq d \leq 120$

b) i) Vertical intercept: 100; horizontal intercept: 4; (0, 100); (4, 0)

ii) -25 km/h

iii) Domain: $0 \leq t \leq 4$; range: $0 \leq d \leq 100$

5. a) i) 400; (0, 400)

ii) 100 ft./min


iii) Domain: $0 \leq t \leq 8$; range: $400 \leq A \leq 1200$

b) i) 1000; (0, 1000)


ii) -50 ft./min

iii) Domain: $0 \leq t \leq 8$; range: $600 \leq A \leq 1000$


6. a)




b)



c)



d)




7. a) $9 \text{ m}^2/\text{L}$; every litre of paint covers an area of 9 m^2 .
 b) 54 m^2
 c) 5 L
8. a) ii
 b) iii
9. a) Vertical intercept: 0; horizontal intercept: 0; (0, 0); the cost of running the backhoe for 0 h is \$0.
 b) \$80/h; each hour that the backhoe is run increases the cost by \$80.
 c) Domain: $0 \leq t \leq 10$; range: $0 \leq C \leq 800$
 d) \$560
 e) 4.5 h
10. a) \$1.50/km; every kilometre driven costs an additional \$1.50.
 b) \$14
 c) 4 km
11. Estimates may vary. Smart car: approximately 0.06 L/km; SUV: approximately 0.128 L/km; the Smart car uses less fuel per kilometre.
12. a) 2.5 h, or 2 h 30 min
 b) 24 km/h
 c) 60 km
 d) $1\frac{2}{3}$ h, or 1 h 40 min
13. a) It takes longer to fill the empty tank.
 b) 25 m^3 of fuel

14. a) Answers may vary. For example:


The scale on the axes is so small that it would be impossible to distinguish every point on the graph.

- b) i) Approximately 33 sweatshirts
 ii) \$15


15. a)




b)



c)



d)



16. a) \$0.80 per bar sold

- b) Vertical intercept: -40; it represents the loss when 0 bars are sold; \$40; horizontal intercept: 50; it represents the number of bars that must be sold to reach the break-even point, when no profit is made and there is no loss: 50 bars
- c) Domain: $0 \leq n \leq 300$, where n is a whole number; range: all multiples of 0.80 from -40 to 200; I wouldn't want to list all the values in the range because there are 301 of them.

17. a) Answers may vary. For example:
There are no intercepts on the graph because the relation does not apply to people less than 10 years of age and older than 90 years of age.

b) Approximately -0.8 (beats/min)/year; for every additional year of age, the recommended maximum heart rate decreases by approximately 1 beat/min.

c) Approximately 77 years of age

d) Approximately 126 beats/min


18. a) i) x -intercept: 5; y -intercept: 5

ii) $x + y = 5$

b) i) x -intercept: 5; y -intercept: -5

ii) $x - y = 5$

19. a)



b) $f(5) = 425$

c) $t = 2.5$

d) Contexts may vary. For example: A car's distance from home as it travels away at an average speed of 85 km/h. In this context, only the 1st quadrant of the graph is relevant.

20. a) The vertical intercept represents the person's distance from Duke Point when starting the journey at Parksville.

The horizontal intercept represents the person's distance from Parksville after completing the journey at Duke Point.

The distance between the two locations doesn't change, so the intercepts have the same value.

b) -1 ; for every 1 km the car moves away from Parksville, it moves 1 km closer to Duke Point.

c) Interchanging the dependent and independent variables would interchange the labels on the axes, but the line on the graph would stay the same.


Chapter 5: Review, page 326

1. a) The table shows a relation with the association "has this cultural heritage" from a set of artists to a set of First Nations heritages.

b) i) As a set of ordered pairs:

$\{(Bob\ Dempsey, Tlingit), (Dorothy\ Grant, Haida), (Bill\ Helin, Tsimshian), (John\ Joseph, Squamish), (Judith\ P.\ Morgan, Gitksan), (Bill\ Reid, Haida), (Susan\ Point, Salish)\}$

ii) As an arrow diagram:




2. Representations may vary. For example:

a) As a table:

Element	Atomic Number
carbon	6
chlorine	17
hydrogen	1
iron	26
oxygen	8
silver	47

As an arrow diagram:




As a set of ordered pairs:

$\{(carbon, 6), (chlorine, 17), (hydrogen, 1), (iron, 26), (oxygen, 8), (silver, 47)\}$

b) As a table:

Atomic Number	Element
1	hydrogen
6	carbon
8	oxygen
17	chlorine
26	iron
47	silver


As an arrow diagram:



As a set of ordered pairs:

$\{(1, \text{hydrogen}), (6, \text{carbon}), (8, \text{oxygen}), (17, \text{chlorine}), (26, \text{iron}), (47, \text{silver})\}$

3. a) Not a function
b) Function
c) Function
d) Not a function
4. a) $f(x) = -4x + 9$
b) $C(n) = 12n + 75$
c) $D(t) = -20t + 150$
d) $P(s) = 4s$
5. a) $P = 5n - 300$
b) Independent variable: n ; dependent variable: P
c) $P(150) = 450$; if 150 students attend the dance, the profit is \$450.
d) $n = 200$; the profit is \$700 when 200 students attend the dance.
6. a) Graph A
b) Answers may vary. For example:
Graph D could represent Laura's journey to school to pick up her bike. She walks to school, then picks up her bicycle and rides home.
7. b) 2 times
c) 2.0 L of water
d) Dependent variable: volume of water in Liam's flask; independent variable: distance Liam hikes
8. a) I joined the points because all times between 0 min and 30 min are permissible and all temperatures between 50°C and 89°C are permissible.



- b) The graph represents a function because a vertical line drawn on the graph passes through one point.

9. Estimates may vary.


- a) Not a function; domain: $\{13, 14, 15, 16, 17\}$; range: $\{159, 161, 165, 168, 170, 174, 176\}$
 - b) Function; domain: $\{08:00, 10:00, 12:00, 14:00, 16:00, 18:00\}$; range: $\{2, 5, 10, 20, 25\}$
10. a) i) Graph A represents the volume of a jar, in cubic centimetres, as a linear function of its height, in centimetres.
ii) Graph B represents the number of marbles in a jar as a linear function of the jar's height, in centimetres.
- b) i) Independent variable: height of the jar, h ;
dependent variable: volume of the jar, V
ii) Independent variable: height of the jar, h ;
dependent variable: number of marbles in the jar, n
 - c) i) Estimates may vary. For example: Domain: $5 \leq h \leq 20$; range: approximately $400 \leq V \leq 1575$
ii) Domain: $\{5, 10, 15, 20\}$; range: $\{14, 28, 42, 56\}$
 - d) The points are joined in Graph A because it is possible for a jar to have any height between 5 cm and 20 cm and any volume between 400 cm^3 and 1575 cm^3 .
The points are not joined in Graph B because only whole numbers of marbles are permissible.

11. a) -2


b) -1

12. Graphs may vary. For example:

a)




b)



13. a) Linear relation b) Linear relation
c) Not a linear relation


14. Tables of values may vary. For example:

a) i)




ii) Table:

x	y
-2	11
-1	5
0	3
1	5
2	11




iii)

x	y
-2	-1
-1	1
0	3
1	5
2	7




iv)




v)

x	y
-2	-6
-1	-3
0	0
1	3
2	6



vi)

x	y
-1	4
0	3
1	2
2	1
3	0



b) i, iii, iv, v, vi

15. a) The equation represents a linear relation because, when g changes by 1, N changes by $\frac{1}{15}$.

b) $\frac{1}{15}$; For every 1 g of carbohydrate that Isabelle consumes, she gives herself $\frac{1}{15}$ of a unit of insulin.

16. a) 6000 m, or 6 km

b) Domain: $0 \leq n \leq 2800$; range: $0 \leq d \leq 6000$

c) Approximately 2.1 m/revolution; in one revolution of the wheel, the bicycle covers a distance of approximately 2 m.

d) Approximately 0.68 m, or 68 cm

17. a) ii

b) iii

c) i

18. a) 201 caps

b) \$4

c) i) 350 caps

ii) 500 caps

d) The profit depends on the sale of caps and the initial cost of \$800 to buy or make the caps. So, doubling the number of caps does not double the profit.

Chapter 5: Practice Test, page 329

1. B


2. C

3. a) i) Function

ii) Representations may vary. For example:

Domain: $\{-3, -1, 0, 1, 2\}$; range: $\{2, 4, 5, 6\}$

As a graph:




The function is not linear because the points on the graph do not lie on a line.

b) i) Function

ii) Representations may vary. For example:

Domain: $\{-3, -1, 1, 2, \dots\}$; range: $\{1, 4, 9, \dots\}$

As a graph:



The function is not linear because the points on the graph do not lie on a line.

c) i) Function

ii) Representations may vary. For example:

Domain: $-2 \leq x \leq 8$; range: $-1 \leq y \leq 4$

As an equation:


$$y = -\frac{1}{2}x + 3, \text{ for } -2 \leq x \leq 8$$

The function is linear because the graph is a non-vertical line.

iii) Independent variable: x ; dependent variable: y ; rate of change: $-\frac{1}{2}$


4. Situation: Jamie's school is 20 km from her home. Jamie rides her friend's bike from school to her friend's home, which is 4 km from her own home. She arrives at her friend's home 20 min after she left school. She talks to her friend for 10 min, then walks the remaining 4 km home in 30 min.

Jamie's Journey Home



5. a) The relation is a function because no number is repeated in the first column.
b) Dependent variable: time; independent variable: mass
c)

Time Needed to Cook a Turkey



I connected the points because both time and mass are not discrete data.

- d) Domain: $4 \leq m \leq 10$; range: $2.5 \leq t \leq 4.0$
e) 0.25 h/kg; for every additional kilogram, the time needed to cook the turkey increases by 0.25 h.
f) 3.25 h or 3 h 15 min

Chapter 6 Linear Functions, page 330

6.1 Slope of a Line, page 339


4. a) $\frac{2}{11}$ b) $\frac{2}{7}$

5. a) Negative
b) Positive
c) Not defined
d) Zero


6. a) Rise: 3; run: 6; slope: $\frac{1}{2}$
b) Rise: -2; run: 8; slope: $-\frac{1}{4}$
c) Rise: 3; run: 4; slope: $\frac{3}{4}$
d) Rise: -6; run: 2; slope: -3
7. a) 3
b) $-\frac{7}{2}$
c) $\frac{1}{2}$
d) $-\frac{1}{2}$

8. Sketches may vary. The lines may be in different positions on the grid but they should have the same orientations as those shown.


a)




b)







c)



d)




9. Sketches may vary. The line segments may have different lengths but they should have the same orientations as those shown.



11. a) $\frac{1}{2}$
 b) $\frac{1}{2}$
 c) The slopes in parts a and b are equal.

12. Diagrams may vary. For example:



b) Similarities: the line segments have the same slope; differences: they pass through different points

13. a) i) 2 ii) $\frac{1}{2}$
 iii) -3 iv) $\frac{1}{3}$

- b) i) As x increases by 1, y increases by 2.
 ii) As x increases by 2, y increases by 1.
 iii) As x increases by 1, y decreases by 3.
 iv) As x increases by 3, y increases by 1.

14. a) Diagrams may vary.

- b) i) The slopes of the segments are equal; all segments on the same line have the same slope.

15. a) $\frac{1}{15}$, or $0.\overline{06}$

b) $13\frac{1}{2}$ in.

16. a) $-\frac{1}{48}$

b) 312 in., or 26 ft.

c) $4\frac{1}{2}$ in.

17. a) Line iv

b) Line iii

c) Line ii

d) Line i

18. a) i) $-\frac{3}{5}$

ii) $\frac{3}{5}$

iii) $-\frac{3}{5}$

iv) $\frac{3}{5}$

b) The slopes of BC and ED are equal. The slopes of BE and CD are equal. The two different slopes are opposites.



19. a) The slope of a horizontal line is 0 because its rise is 0, and the quotient of 0 and any number is zero.

b) The slope of a vertical line is undefined because its run is 0, and the quotient of any number and 0 is undefined; that is, I cannot divide by 0.


20. a) $\frac{1}{3}$

21. Positions of lines on the grid may vary. For example:


a) i)



ii)




iii)



22. 840 cm, or 8.4 m

23. Coordinates may vary.

For example:




- a) $(-4, 5)$, $(-6, -3)$, $(-7, -7)$
 b) $(-7, 3)$, $(-3, -1)$, $(-1, -3)$
 c) $(-8, 2)$, $(-2, 0)$, $(1, -1)$
 d) $(-1, 8)$, $(-9, -6)$, $(-13, -13)$

24. a) i) Positive
 ii) Positive
 iii) Negative
 iv) Not defined


b) Sketches may vary.

For example:



25. a)

Mass and Volume of Aluminum



- b) 2.7 g/cm^3
 c) The slope shows that for every 1 cm^3 increase in the volume of an aluminum cube, the mass of the cube increases by 2.7 g.

d) i) 135 g ii) 742.5 g

e) i) Approximately 37 cm^3

ii) Approximately 167 cm^3

26. a) The number of text messages is restricted to whole numbers.

b) \$0.15, or 15¢ c) \$4.95

d) 48 text messages

e) Assumptions may vary. For example: I assumed that all messages cost the same.

27. a) \$45/month

b) \$505


c) \$55

d) Assumptions may vary. For example: I assumed that Charin continues to save the same amount each month after the 5th month and that the savings account did not earn any interest.

28. a) 2 b) $\frac{2}{3}$


29. No

30. a) Positions of point A may vary. For example:



b) Slope of OB is 2; $\tan \angle AOB = 2$


c)




Slope of OB is $\frac{2}{5}$; $\tan \angle AOB = \frac{2}{5}$

d) The slope of a line segment is equal to the tangent of the angle formed by the segment and the positive x -axis. Both the slope and the tangent are equal to the quotient of the same two numbers.

31. a) The slope is $\tan 30^\circ$, or approximately 0.6.



b) The slope is $\tan 60^\circ$, or approximately 1.7.



c) No

6.2 Slopes of Parallel and Perpendicular Lines, page 349

3. a) $\frac{4}{5}$

b) $-\frac{4}{3}$

c) 3

d) 0

4. a) $-\frac{6}{7}$

b) $\frac{8}{5}$

c) $-\frac{1}{9}$

d) $\frac{1}{5}$

5. a) Parallel

c) Neither

6. a) i) $-\frac{4}{9}$

b) Neither

d) Perpendicular

ii) $\frac{9}{4}$

b) i) 5

ii) $-\frac{1}{5}$

c) i) $\frac{7}{3}$

ii) $-\frac{3}{7}$

d) i) -4

ii) $\frac{1}{4}$

7. Yes; the slope of the line through the golfer's club and the slope of the line through the golfer's feet are the same:

approximately $-\frac{1}{6}$

8. a) i) A(-5, -2), B(1, 5) and C(-1, -4), D(4, 1)

ii) Neither

b) i) E(-3, 4), F(3, 2) and G(2, 5), H(0, -1)

ii) Perpendicular

c) i) J(-2, 3), K(1, -3) and M(3, 1), N(-4, -2)

ii) Neither

d) i) P(0, 5), Q(6, 2) and R(-4, -1), S(0, -3)

ii) Parallel

9. a) Perpendicular

b) Parallel


c) Neither

d) Neither

10. a) Both lines have positive slopes, which are reciprocals.

b) Both lines have positive slopes, which are reciprocals.

11. a) Slope of AB is $\frac{3}{2}$, or 1.5.



b) Slope of CD is $\frac{3}{2}$, or 1.5.

c) Answers may vary. For example:


(1, 2), (3, 5)

d) Slope of AE is $-\frac{2}{3}$.

e) Answers may vary. For example:


(0, -4), (3, -6)

12. a) Slope of AB is -2 .




- b) Slope of CD is -2 .

- c) $(-1, 0)$ and $(0, -2)$



- d) Slope of EF is $\frac{1}{2}$, or 0.5 .

- e) $(4, 0)$ and $(0, -2)$



13. a) Yes

- b) No

14. Trapezoid

15. No

16. The slopes of BC and AC are negative reciprocals, so BC and AC are perpendicular: slope of BC: -2 ; slope of AC: $\frac{1}{2}$

17. Yes; The slopes of DE and EF are negative reciprocals, so DE and EF are perpendicular: slope of DE: $\frac{3}{2}$;

slope of EF: $-\frac{2}{3}$

18. Triangles may vary.

- c) In each case, the line segment that joins the midpoints of two sides of a triangle is parallel to the third side of the triangle.

19. a) No; no pairs of slopes are negative reciprocals.

- b) $D(-2, -1)$

20. Coordinates may vary. For example:

$(3, 7)$, $(-9, 1)$, $(6, 1)$, $(-6, -5)$

21. Rhombuses may vary. The diagonals intersect at right angles.

22. $c = -2$

23. a) $a = 3\frac{1}{4}$, or 3.25

- b) $a = 1\frac{1}{5}$, or 1.2

Chapter 6: Checkpoint 1, page 353

1. Slope of AB: $-\frac{2}{3}$; slope of CD: $\frac{1}{4}$

2. a) $-\frac{15}{4}$

- b) $\frac{5}{3}$

3. Answers may vary. For example: The slope of a line is equal to the slope of any segment of the line, so we can use any two points that form that segment to determine the slope of the line.


4. a) 25 km/h; Jordan's average speed

- b) Approximately 31 km


- c) 2.6 h, or 2 h 36 min

5. The positions of the lines on the grids and their labels may vary. For example:


- a) Neither



- b) Perpendicular



- c) Parallel



6. Coordinates may vary. For example:

- a) $(2, -2)$, $(6, 1)$

- b) $(5, -2)$, $(2, 2)$

7. No, no two of the three slopes of the sides of the triangle are negative reciprocals.
8. Answers may vary. For example: $(-12, 0)$, $(0, -5)$

6.3 Math Lab: Investigating Graphs of Linear Functions, page 356

1. a) From top to bottom:

$$y = \frac{1}{2}x + 4, y = \frac{1}{2}x + 2, y = \frac{1}{2}x - 1,$$


$$y = \frac{1}{2}x - 2, y = \frac{1}{2}x - 3$$

- b) From top to bottom:


$$y = -\frac{1}{3}x + 4, y = -\frac{1}{3}x + 3, y = -\frac{1}{3}x + 1,$$

$$y = -\frac{1}{3}x - 2, y = -\frac{1}{3}x - 3$$

2. m represents the slope and b represents the y -intercept of the line. I could plot the y -intercept, then plot a point using the slope.
3. The graph has a slope of -3 and a y -intercept of 6 .




4. a) All the graphs have y -intercept -1 .
- b)




5. a) All the graphs have slope 1 .


b)




6. a), b)



- c), d)



7. a)



- b) m represents the slope or rate of change; that is, \$15 per person. b represents the initial cost of \$550 to rent the hall.

6.4 Slope-intercept Form of the Equation for a Linear Function, page 362

4. a) Slope: 4 ; y -intercept: -7
 b) Slope: 1 ; y -intercept: 12
 c) Slope: $-\frac{4}{9}$; y -intercept: 7
 d) Slope: 11 ; y -intercept: $-\frac{3}{8}$
 e) Slope: $\frac{1}{5}$; y -intercept: 0
 f) Slope: 0 ; y -intercept: 3

5. a) $y = 7x + 16$

b) $y = -\frac{3}{8}x + 5$


c) $y = \frac{7}{16}x - 3$

d) $y = -\frac{6}{5}x - 8$


e) $y = -\frac{5}{12}x$

6. Sketches may vary. For example:


a)




b)




c)




d)




7. a)




b)




c)




d)



e)



f)




8. a) $C = 50t + 80$


b) $C = 40t + 100$

9. $F = 0.02d + 3.50$


10. a)




b)



c)



d)



11. a) The student may have confused the values of the slope and the y-intercept.

b) $y = 4x - 3$

12. a) i) Slope: $-\frac{1}{2}$; y-intercept: 2

ii) $y = -\frac{1}{2}x + 2$

iii) $y = -3$

b) i) Slope: 4; y-intercept: -6

ii) $y = 4x - 6$

iii) $y = 34$

c) i) Slope: $\frac{3}{4}$; y-intercept: 1

ii) $y = \frac{3}{4}x + 1$

iii) $y = 8.5$

d) i) Slope: $-\frac{1}{3}$; y-intercept: -2

ii) $y = -\frac{1}{3}x - 2$

iii) $y = -\frac{16}{3}$, or $-5\frac{1}{3}$

13. a) Slope: -80; the plane is descending at a speed of 80 m/min. h -intercept: 900; when the plane begins its descent, it is 900 m above the lake.

b) $h = -80t + 900$

c) 460 m

d) i) The graph would be a line joining (0, 700) and (8, 0).

ii) $h = -87.5t + 700$

14. a) $C = 0.80n + 20$

b) \$107.20

c) 125 songs

16. a) $E = 0.05t + 34$

b) \$54

c) \$600

17. a) $y = 4x + 1$

b) $y = \frac{2}{3}x - 1$

c) $y = -\frac{5}{3}x - 7$

18. a) Graph C

c) Graph D

19. a) Graph C

c) Graph B

20. a) Graph B

c) Graph D

b) Graph A

d) Graph B

b) Graph D

d) Graph A

b) Graph C

d) Graph A

21. Parallel lines:

$y = -5x - 7$ and $y = -5x + 13$;

$y = 5x + 15$ and $y = 5x + 24$;

$y = \frac{1}{5}x + 9$ and $y = \frac{1}{5}x + 21$;

$y = -\frac{1}{5}x + 15$ and $y = -\frac{1}{5}x$

Perpendicular lines:

$y = -5x - 7$ and $y = \frac{1}{5}x + 9$;

$y = -5x - 7$ and $y = \frac{1}{5}x + 21$;

$y = -5x + 13$ and $y = \frac{1}{5}x + 9$;

$y = -5x + 13$ and $y = \frac{1}{5}x + 21$;

$y = 5x + 15$ and $y = -\frac{1}{5}x + 15$;

$y = 5x + 15$ and $y = -\frac{1}{5}x$;

$y = 5x + 24$ and $y = -\frac{1}{5}x + 15$;

$y = 5x + 24$ and $y = -\frac{1}{5}x$

22. $y = -\frac{4}{3}x + 4$

23. $c = -\frac{38}{3}$, or $-12\frac{2}{3}$

24. $m = -\frac{47}{24}$, or $-1\frac{23}{24}$


6.5 Slope-Point Form of the Equation for a Linear Function, page 372

4. Coordinates may vary. For example:


- a) Slope: -4 ; $(1, 5)$
- b) Slope: 3 ; $(8, -7)$
- c) Slope: 1 ; $(-15, -11)$
- d) Slope: 5 ; $(2, 0)$
- e) Slope: $\frac{4}{7}$; $(-3, -6)$
- f) Slope: $-\frac{8}{5}$; $(-16, 21)$

- 5. a) $y - 2 = -5(x + 4)$
- b) $y + 8 = 7(x - 6)$
- c) $y + 5 = -\frac{3}{4}(x - 7)$
- d) $y + 8 = 0$, or $y = -8$


6. a)




b)




c)




d)




7. a) The graph is a line through $(4, -2)$ with slope -3 .




b) The graph is a line through $(-3, -4)$ with slope 2 .



c) The graph is a line through $(-5, 3)$ with slope 1 .



d) The graph is a line through $(2, 0)$ with slope -1 .



9. Equations may be written in different forms.

- a) i) $y - 4 = -\frac{4}{3}(x + 2)$ ii) $y - 3 = \frac{2}{5}(x - 3)$
- iii) $y + 2 = \frac{1}{3}(x + 4)$ iv) $y + 2 = -\frac{5}{2}(x - 1)$
- b) i) $y = -\frac{4}{3}x + \frac{4}{3}$; x -intercept: 1 ; y -intercept: $\frac{4}{3}$
- ii) $y = \frac{2}{5}x + \frac{9}{5}$; x -intercept: $-\frac{9}{2}$, or -4.5 ;
 y -intercept: $\frac{9}{5}$
- iii) $y = \frac{1}{3}x - \frac{2}{3}$; x -intercept: 2 ; y -intercept: $-\frac{2}{3}$
- iv) $y = -\frac{5}{2}x + \frac{1}{2}$; x -intercept: $\frac{1}{5}$, or 0.2 ;
 y -intercept: $\frac{1}{2}$, or 0.5

10. Different variables may be used.

a) Let s represent the speed of sound and t represent the air temperature: $s - 337 = 0.6(t - 10)$

b) 331 m/s

11. Slope-point forms of equations may vary. For example:

a) $y - 1 = 2(x - 1)$, or $y + 5 = 2(x + 2)$; $y = 2x - 1$

b) $y + 2 = -(x - 5)$, or $y - 7 = -(x + 4)$; $y = -x + 3$

c) $y - 8 = 3(x - 2)$, or $y + 7 = 3(x + 3)$; $y = 3x + 2$

d) $y + 5 = -2(x + 5)$, or $y + 1 = -2(x + 7)$;
 $y = -2x - 15$

12. a) Graph C: slope 2 and y-intercept -5

b) Graph A: slope 1 and y-intercept 1

c) Graph B: slope 2 and y-intercept 5

d) Graph D: slope -1 and y-intercept -5

13. The graphs are parallel. The graph of $y - y_1 = m(x - x_1)$

passes through the point $P(x_1, y_1)$, and the graph of

$y + y_1 = m(x + x_1)$ passes through the point

$Q(-x_1, -y_1)$.


14. a) $y - 2 = 2(x + 1)$

b) $y - 2 = \frac{1}{3}(x - 1)$


c) $y - 1 = -\frac{2}{3}(x - 2)$

15. Graphs may also be produced on a computer with graphing software. Graphs may show different windows.


a)




b)



c)



d)



16. a) 1.26 g/mL; For every 1 mL of liquid that is poured into the cylinder, the mass of the cylinder and the liquid increases by 1.26 g.

b) Variables and form of the equation may vary.

For example: Let v millilitres represent the volume of the liquid, and M grams represent the mass of the cylinder and liquid; $M - 51.5 = 1.26(v - 20)$

c) 64.1 g

d) 26.3 g

17. a) Variables and form of the equation may vary.

For example: Let M represent the mass of potash in millions of tonnes, and t represent the time in years since 2005; $M = 0.6t + 8.2$

b) 11.2 million tonnes; 14.2 million tonnes;

Assumption: I assume that the relation continues for times beyond 2007 and remains linear.

18. a) Variables and form of the equation may vary. For example: Let p represent the number of students enrolled in francophone schools, and t represent the time, in years, since 2001; $p - 3470 = 198(t - 2)$

b) Approximately 3866 students

19. a) -2 b) $y - 11 = -2(x + 3)$

c) $y + 3 = -2(x - 4)$

20. a) i) $y + 3 = -\frac{4}{3}(x + 5)$

ii) $y + 3 = \frac{3}{4}(x + 5)$

21. a) $y + 2 = 2(x - 1)$ b) $y + 2 = -\frac{1}{2}(x - 1)$

22. a) $y - 6 = -\frac{5}{2}(x - 2)$ b) $y - 6 = \frac{2}{5}(x - 2)$

23. a) $y = \frac{3}{5}(x - 4)$

b) $y + 1 = -\frac{1}{2}(x - 4)$

24. Form of the equation may vary.

$$y = -\frac{9}{2}x + \frac{37}{9}$$


25. Form of the equation may vary.

$$y + 5 = \frac{3}{5}(x + 2)$$

Chapter 6: Checkpoint 2, page 376

1. Screens may vary.

a)



b) Increase the value of m to get a line with a greater slope. Decrease the value of m to get a line with a lesser slope.

c) Increase the value of b to get a line with a greater y -intercept. Decrease the value of b to get a line with a lesser y -intercept.


2. a) Slope: 25; d -intercept: 10; 25 km/h is Eric's average speed; d -intercept: 10 km is Eric's distance from home at the start of his ride.

b) $d = 25t + 10$


c) i) 66.25 km

ii) 1.4 h, or 1 h 24 min


3. a)




b)




c)



d)



e)



4. a) $y = 2x + 3$

b) Equations may have different forms. For example:
 $y - 5 = 2(x - 1)$

6.6 General Form of the Equation for a Linear Relation, page 384

4. a) Standard form

b) General form

c) Slope-intercept form

d) Slope-point form

5. a) x -intercept: 3; y -intercept: -8

b) x -intercept: 8; y -intercept: 7

c) x -intercept: 22; y -intercept: -8

d) x -intercept: 13.5; y -intercept: -3


6. a) $4x + 3y - 36 = 0$

b) $2x - y - 7 = 0$


c) $2x + y - 6 = 0$

d) $5x - y - 1 = 0$

7. a)




b)




8. a) i) The coefficient of x is negative.
 ii) Neither side of the equation is 0.
 iii) The coefficient of x is not a whole number.
 iv) The x -term should come before the y -term.
 b) i) $2x - 3y - 42 = 0$
 ii) $5x - 4y + 100 = 0$
 iii) $x - y + 2 = 0$
 iv) $9x + 5y - 20 = 0$


9. a) i) x -intercept: 8; y -intercept: -6
 ii)



- b) i) x -intercept: -10 ; y -intercept: 12
 ii)




- c) i) x -intercept: 8; y -intercept: -12
 ii)




- d) i) x -intercept: 2; y -intercept: -10

ii)



10. b)




c) $f + s - 12 = 0$

d) Pairs of integers may vary. For example:
 0 and 12; 5 and 7; 3 and 9; 13 and -1 ; 14 and -2 ;
 15 and -3

11. a), b) Letters for the variables may differ.

Let s represent a small pan, and l represent a large pan.

$$12s + 36l = 504$$



12. a) $y = -\frac{4}{3}x + 8$


b) $y = \frac{3}{8}x + \frac{3}{2}$

c) $y = \frac{2}{5}x - 3$


d) $y = -\frac{7}{3}x - \frac{10}{3}$

13. a) -4
 b) 3
 c) 5
 d) -5


14. a)




b)



c)




d)



15. a) 9 pieces of 8-ft. pipe
 b) 12 pieces of 6-ft. pipe
 c) No; 9.75 pieces of 8-ft. pipe would be needed
 d) No; $10\frac{2}{3}$ pieces of 6-ft. pipe would be needed

16. Graphs may have variables on different axes; and variables may be different. Let l represent the number of loonies and t represent the number of toonies.


b), c) $2t + l = 24$




- d) i) No
 ii) No

17. Screens may vary.


a)




b)



c)




d)




18. a) $x - 3y - 12 = 0$
 b) $x - 3y + 11 = 0$
 c) $x + 4y + 11 = 0$
 d) $9x + 6y - 8 = 0$

19. Forms of the equations may vary. For example:


a) $y = \frac{1}{3}x - 4$; $x - 3y - 12 = 0$; $y + 3 = \frac{1}{3}(x - 3)$




b) $y - 2 = \frac{1}{3}(x + 5)$; $x - 3y + 11 = 0$; $y = \frac{1}{3}x + \frac{11}{3}$



c) $y + 3 = -\frac{1}{4}(x - 1)$; $x + 4y + 11 = 0$; $y = -\frac{1}{4}x - \frac{11}{4}$




d) $y = -\frac{3}{2}x + \frac{4}{3}$; $9x + 6y - 8 = 0$; $y + \frac{1}{6} = -\frac{3}{2}(x - 1)$



22. a) Graph B
b) Graph A

23. b)



24. Equations in parts b, e, and g are equivalent.

Equations in parts d, f, and h are equivalent.

26. a) $3x + 4y - 12 = 0$; linear function

b) Not a linear function

c) Not a linear function

d) $x - 3y + 8 = 0$; linear function

28. a) $B \neq 0$: $-\frac{A}{B}$

b) $B \neq 0$: $-\frac{C}{B}$

Chapter 6: Review, page 388

1. a) $-\frac{2}{3}$

b) $\frac{4}{5}$


2. a) Negative

b) Negative

c) Zero


3. Sketches and coordinates may vary.

a) i)




ii) $(-2, 0)$, $(-1, -1)$, $(0, -2)$

b) i)



ii) $(1, 2)$, $(5, 3)$, $(9, 4)$

c) i)



ii) $(-5, 4), (-1, -2), (1, -5)$

4. a) -2

b) $-\frac{3}{2}$

5. a) 160; for every 1 min Gabrielle jogs, she covers a distance of 160 m.

b) Slope is equal to the rate of change.

c) i) 640 m

ii) 6.25 min, or 6 min 15 s

6. a) i) 3 ii) $-\frac{1}{3}$

b) i) $-\frac{6}{5}$ ii) $\frac{5}{6}$

c) i) $\frac{11}{8}$ ii) $-\frac{8}{11}$

d) i) 1 ii) -1

7. a) Perpendicular; slope of JH: 2; slope of KM: $-\frac{1}{2}$

b) Neither; slope of NP: 3; slope of QR: -3


8. No; slope of ST: $-\frac{1}{3}$; slope of TU: 3; slope of UV: $-\frac{4}{9}$;

slope of SV: $\frac{5}{2}$


9. Yes; The slopes of AB and BC are negative reciprocals, so AB and BC are perpendicular.

Slope of AB: 2; slope of BC: $-\frac{1}{2}$

11. a) Slope: -3 ; y-intercept: 4



b) Slope: $\frac{3}{4}$; y-intercept: -2



12. a) i) Slope: $\frac{5}{3}$; y-intercept: 1

ii) $y = \frac{5}{3}x + 1$

b) i) Slope: $-\frac{3}{2}$; y-intercept: -1

ii) $y = -\frac{3}{2}x - 1$

13. a) Graph C

b) Graph D

c) Graph A

d) Graph B

14. a) $A = 15w + 40$

b) 21 weeks

c) The slope would represent the amount Mason saved each week: \$15; the vertical intercept would represent the amount in his bank account when he started saving: \$40

15. Equations may vary. For example:

a) $y = \frac{4}{7}x + 1$ and $y = \frac{4}{7}x - 10$


b) $y = -\frac{7}{4}x + 1$ and $y = -\frac{7}{4}x - 10$

16. $y - 3 = -\frac{1}{2}(x + 2)$

17. Coordinates and forms of the equation may vary.

a) i) 2; $(-3, -4)$


ii)



iii) $y + 2 = 2(x + 2)$

b) i) $-\frac{1}{3}$; $(4, 1)$

ii)



iii) $y - 2 = -\frac{1}{3}(x - 1)$

18. Forms of the equation may vary. For example:

a) $y = \frac{2}{3}(x - 2)$

b) $y - 2 = -\frac{3}{5}(x + 3)$

19. Forms of the equation may vary.

a) i) $y - 5 = 3(x - 1)$ or $y + 7 = 3(x + 3)$

ii) $y + 1 = -\frac{1}{2}(x - 5)$ or $y - 3 = -\frac{1}{2}(x + 3)$

b) Coordinates may vary. For example:

i) (2, 8)

ii) (1, 1)

20. Variables may differ. For example:

a) Let C represent the cost, and p represent the number of people: $C = 44p$

b) \$44

c) 6 people


21. b) i) $5x - 4y + 40 = 0$

ii) $x + 3y - 12 = 0$


iii) $x - 3y + 10 = 0$

iv) $x - 5y + 15 = 0$

22. a) i)




ii)



b) i) $\frac{3}{4}$

ii) $\frac{1}{3}$

24. a), b) $g - l - 6 = 0$



c) Pairs of integers may vary. For example:

8 and 2; 7 and 1; 6 and 0; 5 and -1; 4 and -2

25. Equations in parts a and d are equivalent. Equations in parts b and e are equivalent.


26. a) Graph B

b) Graph C

c) Graph A


27. Variables may differ. Let a represent the number of hours Max babysits for the first family, and b represent the number of hours he babysits for the second family.

a), b) $5a + 4b = 60$



28. Variables may differ. Let n represent the number of new releases and m represent the number of old movies Kylie rents:

a) $5n + 3m = 45$




b) i) No


ii) Yes

Chapter 6: Practice Test, page 391


1. C
2. B
3. a) i)



ii)



iii)



- b) $y - 2 = -\frac{3}{2}(x - 6)$
 - c) $3x + y + 1 = 0$
 - d) Coordinates and equations may vary. For example:
 $P(8, 3)$ and $y = -\frac{2}{7}x + \frac{37}{7}$
4. Answers and forms of equations may vary. For example:
- a) Slope-intercept form: $y = -2x - 2$
 - b) General form: $y + 1 = 0$
 - c) Slope-point form: $y - 1 = \frac{3}{4}(x - 3)$
5. a) \$6570
- b) 520 people

Chapter 7 Systems of Linear Equations

7.1 Developing Systems of Linear Equations, page 401

4. d
5. c
6. a) iii; x dollars represents the cost of a jacket and y dollars represents the cost of a sweater.
- b) i; x represents the length in feet and y represents the width in feet.
- c) ii; x represents the number of chapatti breads sold and y represents the number of naan breads sold.

7. Variables may differ.

a) $2s + 2l = 20$ and $s + 3l = 22$

8. Variables may differ.

a) $2l + s = 24$ and $l - s = 6$

9. a) $3x + y = 17$ and $x = y + 3$

10. $x + 2y = 20$ and $x + y = 13$; Solution B

11. Variables may differ.

$w + j = 60$ and $w - j = 10$; Solution A

15. a) $\frac{C}{B} = \frac{F}{E}$

b) $\frac{C}{A} = \frac{F}{D}$


16. $x + 2y = -8$ and $9x + 10y = 0$

17. a) For example, $3x + 2y = 5$ and $-2x + 3y = 1$


18. b) $x = 3$

7.2 Solving a System of Linear Equations Graphically, page 409


3. a) $x = -4, y = 2$
- b) $x = 2, y = 3$
- c) $x = 1, y = -3$
- d) $x = -2, y = -1$
4. a) $x = 9, y = -2$; exact
- b) $x = -1\frac{3}{4}, y = 2\frac{3}{4}$; approximate
5. a) i) $x = 4, y = 3$




ii) $x = 2, y = 3$



iii) $x = 6, y = -5$



iv) $x = -3, y = 1$




b) The coordinates of the point of intersection represent the solution of the linear system.


6. Approximate

7. Approximations may vary.


a) $x = 2.5, y = -1.5$




b) $x = 1.2, y = 2.2$




c) $x = 3.2, y = 2.5$



d) $x = 3.7, y = -5$




8. a)



b) i) 2500 brochures

ii) It is cheaper to use Company A when fewer than 2500 brochures are printed.


9. a)



b) i) \$30 000

ii) It would be better to choose Plan B when the clerk's monthly sales are less than \$30 000.

10. Variables may differ. $f + r = 391$ and $f - r = 141$




Approximations may vary. For example,

forested area: about 265 hectares;


the rest of the park: about 125 hectares; approximate

11. Variables may differ. $w - l = 43$ and $2w + l = 107$



50 wins and 7 overtime losses; exact


12. Variables may differ. $f + t = 115$ and $5f + 10t = 800$



Seventy \$5 gift cards and forty-five \$10 gift cards; exact

13. Variables may differ.


$$4.8s + 8.0a = 152 \text{ and } s - a = 13$$



7 adults and 20 students; exact


14. Variables may differ.

a) $b + 36g = 1806$ and $b + 24g = 1254$




Approximations may vary. For example, mass of box: 150 g; mass of one golf ball: 46 g; approximate

15. $x = 8\frac{1}{2}$ in. and $y = 12$ in.




16. a) For example, $x = 1.8$ and $y = -0.1$; approximate



17. a) For example, $y = -x + 5$ and $y = -\frac{3}{2}x + 6$

b) $x = 2$ and $y = 3$



18. Equations may vary. For example, $y = -2x - 7$

19. a) The slopes are negative reciprocals: $-\frac{2}{3}$ and $\frac{3}{2}$

b) Answers may vary. For example,

$$y = 4x + 5 \text{ and } y = -\frac{1}{4}x - 2$$


7.3 Math Lab: Using Graphing Technology to Solve a System of Linear Equations, page 412

- Look for equal values of Y_1 and Y_2 , then the corresponding X-value: $x = 4$, $y = 2$
- Graph each line, then determine the coordinates of the point of intersection of the lines.


2. b) $x = 2.\overline{3}$ and $y = -1.1\overline{6}$
 3. 48 cedar tree and 24 spruce tree seedlings
 4. a) i) $x = 1$ and $y = 1$
 ii) $x = 3$ and $y = 0$
 iii) $x = 5$ and $y = -1$
 iv) $x = 7$ and $y = -2$
 b) $x + 2y = 3$ and $2x - y = 21$
 c) $x = 9$ and $y = -3$
 5. No

Chapter 7: Checkpoint 1, page 415

1. Variables may differ.
 a) $2l + 2w = 128$ and $l - w = 16$
 3. $x = 1$ and $y = -1$




4. a)



- b) Plan A is cheaper when the number of visits is greater than 15.
 5. Variables may differ.
 a) $21s + 27a = 396$ and $s + a = 18$

- b) 15 students and 3 adults



6. Variables may differ.
 a) $l + s = 15\,000$ and $1.4l + 0.02s = 7200$
 b) 5000 large trees and 10 000 small trees

7.4 Using a Substitution Strategy to Solve a System of Linear Equations, page 425

4. a) $x = 16, y = -7$ b) $x = 6, y = 7$
 c) $x = -1, y = -8$ d) $x = 1, y = 4$
 5. a) $x = -2, y = 5$ b) $x = -2, y = 3$
 c) $x = 3, y = 5$ d) $x = 1, y = 4$
 6. a) i) $2x, 4x; 4x = 2(2x)$
 ii) $10y, 5y; 10y = 2(5y)$
 iii) $6y, -2y; 6y = -3(-2y)$
 iv) $-3x, 9x; 9x = -3(-3x)$
 b) i) $x = -\frac{1}{2}, y = -1$ ii) $x = 0, y = 1$
 iii) $x = -1, y = 1$ iv) $x = 2, y = 3$
 7. a) i
 b) i) $x = -1, y = 4$ ii) $x = -4, y = 1$
 iii) $x = 5, y = 1$
 8. a) For example, multiply each term in the first equation by 6: $2x - 3y = 12$
 For example, multiply each term in the second equation by 12: $10x + 9y = 12$
 b) $x = 3, y = -2$
 9. a) For example, divide each term in the first equation by 2: $x + y = -2$
 For example, divide each term in the second equation by 4: $-3x + y = -6$
 b) $x = 1, y = -3$

10. Variables may differ.
 $r + n = 186$ and $n - r = 94$
 46 bears responded; 140 bears did not respond.
11. Variables may differ.
 $2l + 2w = 540$ and $l - w = 90$
 Length: 180 cm; width: 90 cm
12. Variables may differ.
 $s + a = 45$ and $0.8s + 0.6a = 31$
 20 students and 25 adults
13. Variables may differ.
 $x + y = 11$ and $4x + 5y = 47$
 8 groups of 4 and 3 groups of 5
14. Variables may differ.
 $p + a = 85$ and $0.6p + 0.4a = 38$
 20 people masks; 65 animal masks
15. Variables may differ.
 $0.80A + 0.92B = 63$ and $A + B = 75$
 Part A: 50 marks; part B: 25 marks
16. Variables may differ.
 $x + y = 5000$ and $0.025x + 0.0375y = 162.50$
 Two thousand dollars in the 2.5% bond; \$3000 in the 3.75% bond
17. Variables may differ.
 $76s + 49d = 474.25$ and $54s + 37d = 346.25$
 Single-scoop cone: \$3.50; double-scoop cone: \$4.25
18. Joel would have to work 15 weekends before he earns the same amount as Sue.
19. a) $x = 6, y = -3$ b) $x = -1, y = \frac{1}{3}$
 c) $x = -\frac{42}{13}, y = -\frac{72}{13}$ d) $x = \frac{124}{51}, y = -\frac{16}{17}$
20. b) $r = 20, c = 5$
21. $x = 5, y = 22$
22. a) For example: $4x - 2y = -8$ and $9x + 6y = 3$
 b) $x = -1, y = 2$; the systems have the same solution.
23. a) 16 km/h
 b) 40 km
24. Mean mass of males: 205.7 g; mean mass of females: 168 g
25. Rate of climb: 200 m/min; rate of descent: -200 m/min
27. $A = 4, B = -3$

7.5 Using an Elimination Strategy to Solve a System of Linear Equations, page 437

3. a) $x = -3, y = -1$ b) $a = \frac{5}{3}, b = 0$
 c) $x = -1, y = -1$ d) $x = 4, y = 3$
4. a) i) $3x - 6y = -18$ and $3x - y = 2$
 ii) $x - 2y = -6$ and $6x - 2y = 4$
 b) i) $15x - 2y = 9$ and $15x + 12y = 51$
 ii) $-30x + 4y = -18$ and $5x + 4y = 17$
 c) i) $35x + 15y = 45$ and $35x + 14y = 49$
 ii) $14x + 6y = 18$ and $15x + 6y = 21$

- d) i) $42x + 45y = 48$ and $42x + 20y = -2$
 ii) $28x + 30y = 32$ and $63x + 30y = -3$
5. a) $x = 2, y = 4$ b) $x = 1, y = 3$
 c) $x = 3, y = -4$ d) $x = -1, y = 2$
6. a) $x = -4, y = 3$ b) $m = -\frac{2}{3}, n = -\frac{1}{3}$
 c) $s = 0, t = 2$ d) $a = 3, b = -2$
7. a) $x = \frac{79}{7}, y = \frac{122}{7}$ b) $a = -3, b = -7$
 c) $a = \frac{1}{2}, b = \frac{1}{3}$ d) $x = \frac{5}{2}, y = -3$
8. Variables may differ.
 $x + y = 90$ 530 and $y - x = 120$
 2006 attendance: 45 205; 2008 attendance: 45 325
9. Variables may differ.
 $t + s = 545$ and $t - s = 185$
 Talise's dress: 365 cones; her sister's dress: 180 cones
10. Variables may differ.
 $10k + 20b = 200$ and $15k + 25b = 270$
 1 knife: 8 beaver pelts; 1 blanket: 6 beaver pelts
11. Variables may differ.
 $4.5m + 0.5f = 620$ and $f - m = 40$
 Moderate tempo: 120 beats/min; fast tempo: 160 beats/min
12. a) $a = \frac{4}{5}, b = \frac{9}{5}$ b) $x = 20, y = -6$
 c) $x = -0.35, y = 0.25$ d) $x = 0.5, y = 0.5$
13. 18 Canadian; 7 foreign
14. 36 girls; 40 boys
15. a) $3x + y = 17$ and $x + y = 7$
 b) From Balance scales 2, the sum of mass x and mass y is 7 kg. The same mass is being removed from each pan. So, the scales will still be balanced.
 c) Two x -masses equal 10 kg. So, mass x is 5 kg. Remove mass x from the left side of Balance scales 2 and 5 kg from the right side. Then mass y balances 2 kg.
 d) When I remove the x mass, y mass, and 7 kg from Balance scales 1, it is like subtracting the second equation from the first equation to eliminate y .
16. An adult pays \$6.75 and a child pays \$7. So, a child's ticket is more expensive.
17. 15 kg of green peas; 10 kg of red lentils
18. Problems may vary. $x = 5, y = 3$
19. b) $x = 5, y = 2$
20. a) For example, multiply equation 1 by -2 and equation 2 by 3, then add to eliminate x . Multiply equation 1 by 5 and equation 2 by 4, then add to eliminate y .
 b) $x = 3, y = 5$
22. \$950 in the stock; \$450 in the bond
23. a) For example, $3x + 6y = 9$; $x = -1, y = 2$

- b) The solution to each system is: $x = -1, y = 2$
 c) The solutions are the same.
24. a) 40 bushels/acre for wheat; 58 bushels/acre for barley
 b) No, I could use the solution to part a and proportions to determine the yield in bushels/hectare.

Chapter 7: Checkpoint 2, page 441

1. a) $x = \frac{1}{2}, y = \frac{3}{2}$ b) $x = 0, y = -1$
 c) $x = -6, y = -1$
 2. a) Variables may differ.
 $6x + 7y = 494$ and $x - y = 13$
 b) 45 replicas with 6 stones; 32 replicas with 7 stones
 3. \$500 was invested in each bond.
 4. a) $x = -6, y = -7$ b) $x = \frac{1}{2}, y = 3$
 c) $x = -0.75, y = -1.75$ d) $x = -\frac{14}{5}, y = \frac{2}{5}$
 5. Soup: 90 times; a main course: 70 times
 6. Larger volume: 1450 mL; smaller volume: 450 mL
 7. $x = 55^\circ; y = 65^\circ$


7.6 Properties of Systems of Linear Equations, page 448

4. a) i) 1 ii) -1
 iii) 1 iv) -1
 b) i and iii; ii and iv
 c) i and ii; i and iv; ii and iii; iii and iv
 5. a) A and C; B and C b) A and B
 6. a) For example, $x - 3y = 12$ and $5x - 15y = -60$
 b) For example, $6x + 3y = 5$ and $2x - 6y = 24$
 c) For example, $4x + 2y = 20$ and $2x + y = 10$
 7. a) One solution
 b) Infinite solutions
 c) No solution d) No solution
 8. a) For example, $y = x + 2$
 b) For example, $y = 2x + 2$
 c) For example, $-4x + 2y = 2$
 9. a) No solution b) One solution
 c) One solution
 10. One solution
 11. I need to know whether the y -intercepts are the same or different.
 12. For example:
 One solution: $-3x - 4y = 12$
 No solution: $3x - 4y = 8$
 Infinite solutions: $6x - 8y = 24$
 13. Infinite solutions
 14. One solution
 15. Infinite solutions
 16. One solution
 17. One solution


18. 0 points of intersection: slopes of the lines are equal and their y -intercepts are different.
 1 point of intersection: slopes of the lines are different.
 Infinite points of intersection: slopes of the lines are equal and their y -intercepts are equal.
19. a) For example: $x + y = 5$ and $2x + 2y = 10$
 b) When I try to eliminate one variable, I also eliminate the other variable and the constant.
20. a) For example: $x + y = 4$ and $2x + 2y = 6$
 b) When I try to eliminate one variable, I also eliminate the other variable.
22. a) i) Infinite solutions
 ii) No solution iii) One solution
24. a) i) $k \neq \frac{3}{4}$ ii) $k = \frac{3}{4}$

Chapter 7: Review, page 452

1. a) Variables may differ.
 $o + s = 41$ and $o - s = 17$
 b) Solution B
2. a) Variables may differ.
 $s + l = 25$ and $15s + 25l = 475$
 b) Solution B
4. a) $3x + y = 11$ and $3x - 5y = -1$
 b) $x = 3, y = 2$; exact
5. a) George: draw a line through each pair of points, then determine the coordinates of the point of intersection.
 Sunita: plot each y -intercept, then use the slope to mark another point on each line.



7. a)




The graphs appear to intersect at $(-2.8, -6.1)$.

- b) Exact; when $(-2.8, -6.1)$ is substituted into each equation, the left side equals the right side.
8. a) Variables may differ.
 $2c + 4b = 940$ and $c + 3b = 620$

- b) Each line represents one of the equations in the linear system.
- c) One bowl of cereal has 170 mg of sodium and 1 slice of bacon has 150 mg of sodium; exact solution.
9. Where necessary, the answers have been written to 3 decimal places.
- a) $x \doteq 1.526$, $y \doteq 3.316$ b) $x = 12$, $y = 0$
- c) $x = 3.25$, $y = -1.4$
- d) $x \doteq -6.071$, $y \doteq 1.964$
10. a) $x = 0$, $y = -5$ b) $x = 1$, $y = 3$
- c) $x = \frac{19}{7}$, $y = -\frac{11}{63}$ d) $x = -1$, $y = -2$
11. c) $x = -1$, $y = 8$
12. a) Variables may differ.
 $\frac{1}{4}x + \frac{2}{3}y = 5\frac{3}{4}$ and $x - y = 1$
- b) 7 one-quarter cup measures; 6 two-third cup measures

13. a)



- b) Variables may differ.
 $60l + 2w = 306$ and $2l + 60w = 190$
- c) Width: 3 ft.; length: 5 ft.
14. 35 triangles; 115 squares
15. a) $x = 0$, $y = -5$ b) $x = -\frac{11}{2}$, $y = -6$
16. c) $x = 2.5$, $y = -0.25$
17. a) $2l + \left(1 + \frac{1}{2}\pi\right)w = 68\frac{5}{6}$ and $l - w = 7$
- b) Length: 19 ft.; width: 12 ft.
18. a) Infinite solutions, for example:
 $x + y = -1$ and $2x + 2y = -2$
 No solution, for example: $2x + 2y = 5$ and $4x + 4y = -5$
19. a) Clue 1 and Clue 2 b) 45 and 12
20. a) No solution
- b) Infinite solutions
- c) One solution d) No solution

Chapter 7: Practice Test, page 455


1. B
2. A

4. b) $s = 6$, $a = 2$
5. a) i) $x = -4$, $y = \frac{7}{2}$ ii) $x = 4$, $y = 5$
- iii) $x = \frac{3}{2}$, $y = \frac{1}{2}$
- b) The solution of a linear system is the coordinates of the point of intersection of the graphs of the lines.
6. a) Variables may differ.
 $y + r = 90$ and $25y + 12.5r = 1500$
- b) 30 squares and 60 triangles


Cumulative Review Chapters 1–7, page 458

1. Answers may vary. These answers are calculated using exact conversions.
- a) 9 ft. 6 in. b) 457 cm
- c) 4 mi. 1709 yd. d) 165 m
- e) 269 ft. f) 25.75 km
2. a) 384 cm^2 ; 384 cm^3 b) 579 in.^2 ; 924 in.^3
- c) 254 cm^2 ; 382 cm^3
3. 56.3°
4. $36\frac{4}{10}$ in.
5. a) $81 + 18s + s^2$ b) $6a^2 - 19a + 15$
- c) $10n^2 + 7np - 12p^2$ d) $64s^2 - t^2$
- e) $-2w^3 - w^2 + 20w - 32$
- f) $-6x^4 + 5x^3 + 22x^2 + 2x - 8$
6. a) $7(2a^3b^2 - 4b^3c^2 + 3a^2c^3)$
- b) $(n - 4)(n + 3)$
- c) $4(3r + 4s)(3r - 4s)$ d) $(2m + 9)(3m - 2)$
- e) $(w - 11x)^2$ f) $(5c + 6d)(6c - 5d)$
7. a) i) $3\sqrt{5}$ ii) $4\sqrt[3]{2}$
- iii) $\sqrt[4]{932}$ iv) $7\sqrt{11}$
- b) i) $\sqrt{432}$ ii) $\sqrt[3]{189}$
- iii) $\sqrt[5]{480}$ iv) $\sqrt{425}$
8. a) $\frac{a^2}{b^5}$ b) $\frac{c^2}{d^5}$
- c) $-\frac{x^5}{4y^3z^4}$ d) $-\frac{6}{a^3b^2}$
9. a) $\frac{9}{16}$ b) 12.25
- c) $\frac{25}{36}$ d) 2.5
10. a) The relation shows the association “participates in” from a set of athletes to a set of sports.
- b) i) {Perdita Felicien, track), (Donovan Bailey, track), (Nancy Greene, skiing), (Annamay Pierse, swimming), (Justin Morneau, baseball), (Steve Nash, basketball)}

ii)




11. a) Each number in the first column of the table appears exactly once.
 b) Independent variable: v ; dependent variable: C
 c) Domain: $\{1, 2, 3, 4, \dots\}$; range: $\{1.09, 2.18, 3.27, 4.36, \dots\}$
 d) $C(v) = 1.09v$
 e) $C(25) = 27.25$; the cost of 25 L of gasoline is \$27.25.
 f) $v \doteq 46$; with \$50, approximately 46 L of gasoline can be purchased.
12. a) False b) True
 c) True d) False
13. a) Graph B
14. a) Domain: $x \leq 3$; range: $y \geq -2$
 b) Domain: all real numbers; range: $y \leq 3$
15. a) Tables of values and sketches may vary. For example:




ii)

x	y
-2	5
-1	3
0	1
1	-1
2	-3




iii)

x	y
-2	2
-1	-1
0	-2
1	-1
2	2




iv) A vertical line that passes through $(2, 0)$




v)

x	y
-2	4
0	3
2	2
4	1
6	0




vi)


x	y
1	0
2	1
2	-1
5	2
5	-2




- b) i, ii, iv, and v; the graphs are straight lines.
16. a) 300; $(0, 300)$; the fixed cost of renting the banquet room is \$300.
 b) \$15/person; for each additional person who attends, the cost increases by \$15.
 c) Domain: $\{0, 1, 2, 3, 4, 5, \dots\}$; range: $\{300, 315, 330, 345, 360, 375, \dots\}$
 The domain can be any whole number up to the number of people the banquet room can hold.
 The range can be any multiple of 15 greater than or equal to 300, up to a number that depends on the maximum capacity of the room.
 d) \$1050 e) 25 people
17. Points and sketches may vary. For example:
- a) $(-3, -1)$, $(-2, 1)$, $(0, 5)$




- b) $(-2, 6), (0, 0), (1, -3)$




- c) $(-5, 0), (3, 6), (7, 9)$



- d) $(-6, 5), (4, 1), (9, -1)$



18. a) Neither
b) Perpendicular
c) Parallel
d) Perpendicular
19. a)



- b) Changing the value of m changes the steepness of the graph.

20. a) The student wrote the slope as $\frac{1}{2}$ instead of -2 , and the y -intercept as -3 instead of 3 .

- b) $y = -2x + 3$

21. a) Equations may vary. For example:

i) $y - 2 = -\frac{4}{5}(x + 1)$

ii) $y + 3 = 2(x + 2)$


- b) i) $y = -\frac{4}{5}x + \frac{6}{5}$; x -intercept: $\frac{3}{2}$; y -intercept: $\frac{6}{5}$

ii) $y = 2x + 1$; x -intercept: $-\frac{1}{2}$; y -intercept: 1


22. a) $d = 14t + 200$
b) \$690
c) 37 h
d) No

23. a) Sketches may vary. For example:

i)



ii)



- b) i) -7
ii) $\frac{2}{5}$

24. a) i) $25x - 20y - 12 = 0$

ii) $2x - 3y - 14 = 0$

25. Variables may differ.

$$9.60s + 20.80l = 2206.40 \text{ and } s + l = 140$$

26. a) Forms of equations in the system may vary.


For example:

$$8a + 5s = 1122 \text{ and } s - a = 32$$

- b) Answers may vary. For example:

75 adults and 105 students; approximate

(Actual answer: 74 adults and 106 students)



27. $x = \frac{8}{3}$; $y = \frac{7}{12}$

28. Part A: 48 marks; Part B: 60 marks

29. a) $x = \frac{53}{26}$, $y = -\frac{8}{13}$

b) $x = -3$, $y = \frac{5}{2}$

30. Equations may vary. For example:

One solution: $x - y = 1$

No solution: $5x + 3y = 1$

Infinite solutions: $10x + 6y = 30$